

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/099092 A2

- (51) International Patent Classification⁷: C12N 9/00
- (21) International Application Number: PCT/DK02/00388
- (22) International Filing Date: 7 June 2002 (07.06.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/296,142 7 June 2001 (07.06.2001) US
- (71) Applicant (for all designated States except US): HEME-BIOTECH A/S [DK/DK]; Roskildevej 12C, DK-3400 Hillerød (DK).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): FOGH, Jens [DK/DK]; Bjergagervej 37, DK-3540 Lyngby (DK). IRANI, Meher [US/US]; 3503 N.E. 100th, Seattle, WA 98125 (US). ANDERSSON, Claes [SE/SE]; Tallåvågen 5, S-187 43 Täby (SE). WEIGELT, Cecilia [SE/SE]; Banérsgatan 27, 1 tr., S-115 22 Stockholm (SE).
- (74) Agent: PLOUGMANN & VINGTOFT A/S; Sundkrogs-gade 9, P.O. Box 831, DK-2100 Copenhagen Ø (DK).
- (81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
- without international search report and to be republished upon receipt of that report
 - with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/099092 A2

(54) Title: PRODUCTION OF RECOMBINANT HUMAN LYSOSOMAL ALPHA-MANNOSIDASE

(57) Abstract: The present invention relates to a cell capable of producing recombinant human LAMAN (rhLAMAN), said cell comprising the 3066 basepair EcoRI - XbaI fragment of a human cDNA which codes for a human LAMAN protein in which the position corresponding to position 186 of the full length hLAMAN protein is Aspartic acid (Asp, D). In particular, the invention relates to a cell comprising a DNA sequence which codes for the amino acid sequence shown in SEQ ID NO. 12. One embodiment is an expression plasmid pLamanExp1 having the sequence shown in SEQ ID NO. 2. One aspect of the invention relates to a method for the preparation of recombinant human LAMAN, the method comprising a) introducing, into a suitable vector, a nucleic acid fragment comprising a DNA fragment which codes for the amino acid sequence shown in SEQ ID NO. 12, b) transforming a cell with the vector obtained in step a), c) culturing the transformed host cell under conditions facilitating expression of the nucleic acid sequence, d) recovering the expression product from the culture. The method may further comprise a fermentation step and/or a purification step. The invention further relates to a rhLAMAN produced by the method of the invention and to the use of the rhLAMAN produced according to the invention for the preparation of a medicament for the treatment of alfa mannosidosis.

WO 02/099092

PCT/DK02/00388

1

Production of recombinant human lysosomal alpha-mannosidase

- The present invention relates to a cell capable of producing recombinant human LAMAN (rhLAMAN), said cell comprising the 3066 basepair EcoRI - XbaI fragment of a human
- 5 cDNA which codes for a human LAMAN protein in which the position corresponding to position 186 of the full length hLAMAN protein is Aspartic acid (Asp, D). In particular, the method relates to a cell comprising a DNA sequence which codes for the amino acid sequence shown in SEQ ID NO 12.
- 10 In a further aspect, the invention relates to a method for the preparation of recombinant human LAMAN, the method comprising a) Introducing, into a suitable vector, a nucleic acid fragment comprising a DNA fragment which codes for the amino acid sequence shown SEQ ID NO 12; b) transforming a cell with the vector obtained in step a); c) culturing the transformed host cell under conditions facilitating expression of the nucleic acid sequence;
- 15 d) recovering the expression product from the culture. The method may further comprise a fermentation step and/or a purification step.

- Another embodiment of the present invention relates to a method for preventing or treating the development of symptoms related to alpha-mannosidosis caused by a de-
- 20 ficiency, in a patient, of the human lysosomal alpha-mannosidase (hLAMAN) enzyme, the method comprising administering over cellular membranes, to a target cell, an effective amount of recombinant human Lysosomal alpha-mannosidase (rhLAMAN) enzyme.

BACKGROUND OF THE INVENTION

25

Alpha-mannosidosis

- Alpha-mannosidosis is a recessive autosomal disease that occurs world wide with a frequency of between 1/1.000.000 and 1/500.000. Mannosidosis is found in all ethnic
- 30 groups in Europe, America, Africa and also Asia. It is detected in all countries with a good diagnostic service for lysosomal storage disorders, at a similar frequency. They are born apparently healthy, however the symptoms of the diseases are progressive. Alpha-mannosidosis displays clinical heterogeneity, ranging from very serious to very mild forms. Typical clinical symptoms are: mental retardation, skeletal changes, impaired immune
- 35 system resulting in recurrent infections, hearing impairment and often the disease is associated with a typical facial characteristics such as a coarse face, a prominent forehead, a flattened nasal bridge, a small nose, and a broad mouth. In the most severe cases (mannosidosis type I) the children suffer from hepatosplenomegaly, and they die during

WO 02/099092

PCT/DK02/00388

2

the first years of life. Possibly this early death is caused by severe infections due to the immunodeficiency caused by the disease. In milder cases (mannosidosis type 2) the patients usually reach adult age. The skeletal weaknesses of the result in the needs of wheeling chairs at age 20 to 40. The disease causes a diffuse dysfunction of the brain
5 often resulting in weak mental performances that excludes anything but the most basic skills of simple reading and writing. These problems associated with hearing inabilities and other clinical manifestations preclude the patient from an independent life, the consequence being that life long caretaking is needed.

10 Lysosomal alpha-mannosidase

Alpha-mannosidosis results from a deficient activity of lysosomal alpha-mannosidase (LAMAN, EC3.2.1.24). The disease is characterised by massive intracellular accumulation of mannose-rich oligosaccharides. LAMAN is an exoglycosidase which hydrolyses terminal, non-reducing alpha-D-mannose residues in alpha-D-mannosides from the non-reducing
15 end during the ordered degradation of the N-linked glycoproteins (Aronson and Kuranda FASEB J 3:2615-2622. 1989). The human enzyme is synthesised as a single polypeptide of 1011 amino acids with a putative signal peptide of 49 residues that is processed into three main glycopeptides of 15, 42, and 70 kD. (Nilssen et al. Hum.Mol.Genet. 6, 717-726. 1997).

20

The lysosomal alpha-mannosidase gene

The gene coding for LAMAN (MANB) is located at chromosome 19 (19cen-q12), (Kaneda et al. Chromosoma 95:8-12. 1987). MANB consists of 24 exons, spanning 21.5 kb (GenBank accession numbers U60885-U60899; Riise et al. Genomics 42:200-207 . 1997). The
25 LAMAN transcript is » 3,500 nucleotides (nts) and contains an open reading frame encoding 1,011 amino acids (GenBank U60266.1).

The cloning and sequencing of the human cDNA encoding LAMAN has been published in three papers (Nilssen et al. Hum.Mol.Genet. 6, 717-726. 1997; Liao et al. J.Biol.Chem.
30 271, 28348-28358. 1996; Nebes et al. Biochem.Biophys.Res.Comm. 200, 239-245. 1994). Curiously, the three sequences are not identical. When compared to the sequence of Nilssen et al (accession # U60266.1) a TA to AT change at positions 1670 and 1671 resulting in a valine to aspartic acid substitution was found by Liao et al. and Nebes et al.

35 Mutations

In 1997 Nilssen et al. (Hum Mol Genet 6:717-726) reported the first disease causing mutation in the fibroblasts from two siblings with mannosidosis in a highly inbred Palestinian family. In these two patients, the cause of the disease is a point mutation at the gene, with the nucleotide Adenine substituted with Thymidin. Since then, several

WO 02/099092

PCT/DK02/00388

3

genetic causes to the disease have been found in man. Today mutations causing alpha-mannosidosis have been characterised in more than 60 patients, mainly in Europeans. These mutations vary both in type, encompassing nucleotide substitutions, deletions and additions and spanning most of the 24 exons of the gene (Berg et al. Am J Genet 64: 77-88. 1999).

The consequences of the mutations on the protein level also vary, from amino acid substitutions that effect both the active site and the folding, to early frame shifts and premature stop codons, resulting in a completely inactive product.

10

The amino acid sequence and enzyme structure in a number of different species. When comparing species, many of the human and the bovine mutations found are located to a highly "conserved" region of the mannosidase polypeptide for several species like man, cattle, cat, whale, bird, cod and slime mould, indicating that these regions are very important for the enzyme function.

Despite the many different types of mutations, it seems that all the mutations cause the similarly complete loss of enzyme activity in cells. Thus, apparently, the variations of clinical severity's cannot be explained by variations in the enzyme activity. Rather, these variations are caused by other factors, being attributed to the general ability of the organism to buffer the obnoxious effects of the accumulating oligomannosides, and by the environments to prevent and to treat infections and other clinical symptoms in the patients. Since no patients appear to possess enzyme activity, alpha-mannosidosis may be regarded as a mild disease compared to many other lysosomal storage disorders, but similar to the other glycoproteinoses (aspartylglucosaminuria, fucosidosis, Schindler's disease, beta-mannosidosis, Salla's disease).

Based on this one may assume that even a very small level of enzyme activity will be sufficient for a normal phenotype within this category of diseases.

30

Diagnosis

The diagnosis of alpha-mannosidosis is currently is based on clinical evaluation, detection of mannose-rich oligosaccharides in urine, and direct measurements of alpha-mannosidase activity in various cell types, such as leukocytes, fibroblasts, and amniocytes (Chester et al., In: Durand P, O'Brian J (eds) Genetic errors of glycoprotein metabolism. Edi-Ermes, Milan, pp 89-120. 1982; Thomas and Beaudet . In: Scriver CR, Beaudet AL, Sly WA, Valle

WO 02/099092

PCT/DK02/00388

4

D (eds) The metabolic and molecular bases of inherited disease. Vol 5. McGraw-Hill, New York, pp 2529-2562. 1995).

Because the symptoms initially often are mild and the biochemical diagnosis is difficult, the
5 diagnosis is frequently made late in the course of the disease. It is obvious that patients
and their families would benefit substantially from an early diagnosis.

Animal models

Also animals may develop alpha-mannosidosis.

10

Alpha mannosidosis has been described in cattle (Hocking et al. Biochem J 128:69-78.
1972), cats (Walkley et al. Proc. Nat. Acad. Sci. 91: 2970-2974, 1994), and guinea pigs
(Crawley et al. Pediatr Res 46: 501-509, 1999). A mouse model was recently generated by
targeted disruption of the alpha-mannosidase gene (Stinchi et al. Hum Mol Genet 8: 1366-
15 72, 1999).

Like in humans alpha mannosidase seems to be caused by specific mutations in the gene
coding for lysosomal alpha-mannosidase. Berg et al. (Biochem J. 328:863-870.1997)
reported the purification of feline liver lysosomal alpha-mannosidase and determination of
20 its cDNA sequence. The active enzyme consists of 3 polypeptides, with molecular masses
of 72, 41, and 12 kD. Similary to the human enzyme it was demonstrated that the feline
enzyme is synthesized as a single-chain precursor with a putative signal peptide of 50
amino acids followed by a polypeptide chain of 957 amino acids, which is cleaved into the
3 polypeptides of the mature enzyme. The deduced amino acid sequence was 81.1% and
25 83.2% identical with the human and bovine sequences, respectively. A 4-bp deletion was
identified in an affected Persian cat; the deletion resulted in a frameshift from codon 583
and premature termination at codon 645. No enzyme activity could be detected in the liver
of the cat. A domestic long-haired cat expressing a milder phenotype had enzyme activity
of 2% of normal; this cat did not possess the 4-bp deletion. Tollersrud et al. (Eur J
30 Biochem 246:410-419 .1997) purified the bovine kidney enzyme to homogeneity and
cloned the gene. The gene was organized in 24 exons that spanned 16 kb. Based on the
gene sequence they identified two mutations in cattle.

35 Medical need for alpha-mannosidosis therapy

The lack of effective treatment for alpha-mannosidosis is well recognised.

WO 02/099092

PCT/DK02/00388

5

Enzyme substitution

When lysosomal storage diseases were discovered, hopes were raised that this could be treated by enzyme substitution. Enzyme replacement therapy has proven efficient in Gaucher disease. When exogenous lysosomal glucocerebrosidase is injected into the

5 patient, this enzyme is taken up by enzyme-deficient cells (Barton et al. N Engl J Med 324:1464-1470). Such uptake is regulated by certain receptors on the cell surface. Different tissue have different receptors. It is therefore necessary to change the natural enzyme to enable uptake into the tissue of interest. If properly designed, the missing enzyme could be injected regularly like the diabetic subject injects insulin. In many In vitro

10 studies with the purified active enzyme added to the media of enzyme-deficient fibroblasts, the lysosomal substrate accumulation was corrected, but In vivo treatment has been hampered through the problem of producing the sufficient quantity of enzymes, immunological problems, and that the substituted enzyme does not pass the blood-brain barrier.

15

Bone marrow transplantation

In 1996 Walkley et al. (Proc. Nat. Acad. Sci. 91: 2970-2974, 1994) published a paper on 3 kittens with mannosidosis that were treated with bone marrow transplantation (BMT) in 1991. In the 2 animals that were sacrificed a normalisation was seen, not only in the body,

20 but more importantly, also in brain. The 3rd cat was well after 6 years. Normally, an untreated cat dies with 3-6 months. In 1987 a child with mannosidosis was treated with BMT (Will et al. Arch Dis Child 1987 Oct;62(10):1044-9). He died after 18 weeks due to procedure related complications. In brain little enzyme activity was found. This disappointing result could be explained by heavy immunosuppressive treatment before

25 death, or that it takes time for the enzyme activity to increase in brain after BMT. The donor was the mother (who as carrier must be expected to have less than 50% enzyme activity) or it may be BMT in man has no effect on enzyme function in brain.

DISCLOSURE OF THE INVENTION

30

A particularly important embodiment of the present invention is a cell capable of producing recombinant human LAMAN (rhLAMAN), said cell comprising the approximately 3066 basepair EcoRI - XbaI fragment of a human cDNA which codes for a human LAMAN protein in which the position corresponding to position 186 of the full length hLAMAN protein is

35 Aspartic acid (Asp, D). In a recent paper Berg et al. (Mol Gen and Metabol 73, 18-29, 2001) described a cell capable of producing rhLAMAN. Whereas Berg et al. does not disclose the sequence of the particular cDNA used in the paper the same group has published their LAMAN sequence as GenBank accession no. U60266. In this sequence the amino acid corresponding to position 186 is Valine not Aspartic acid as in the present

WO 02/099092

PCT/DK02/00388

6

application. Interesting, Nebes et al. (Biochem.Biophys.Res.Commun. 200, 239-245. 1994) reports that the amino acid at position 186 is Aspartic acid. However, the sequence of Nebes et al. differs from the sequence disclosed herein at many other positions. For instance the position corresponding to position 343 in the amino acid sequence is missing
5 in Nebes et al., also the start of the protein sequence disclosed herein is absent in Nebes et al.

A further preferred embodiment of the present invention is a cell which contains a DNA fragment comprising the approximately 3066 basepair EcoRI - XbaI fragment of the DNA
10 fragment having the sequence shown in SEQ ID NO 2. In addition to coding for full length hLAMAN the sequence is engineered to comprise two convenient restriction enzyme sites for easy insertion into a wide range of suitable expression vectors.

A most preferred embodiment of the present invention is a cell obtained by use of the
15 expression plasmid pLamanExp1 since this plasmid contains signal sequences necessary to obtain high expression levels and such a cell furthermore is suitable for amplification as described in example 2 .

Another inventive concept of the present invention is based on the novel idea of
20 substituting the reduced hLAMAN enzymatic activity in the person having alpha-mannosidosis simply by administering over cellular membranes, to a target cell, an effective amount of an hLAMAN enzyme. This can thereby "assist" the enzyme that is in deficit.

25 Accordingly, the present invention relates to a method for preventing or treating the development of symptoms related to alpha-mannosidosis(alpha-mannosidosis) caused by a deficiency, in a subject, of the human Lysosomal alpha-mannosidase (hLAMAN) enzyme, the method comprising administering over cellular membranes, to a target cell, an effective amount of an hLAMAN catalyst which is said enzyme or an enzymatically
30 equivalent part or analogue thereof.

Since mannose-rich glycoproteins accumulates in all cells of the body and causes the symptoms of mannosidase, the target cells for hLAMAN therapy are all cells of the body. Since the some of the most severe symptoms caused by hLAMAN deficiency are related to
35 the central nerve system (CNS) the CNS including the brain is the one important objective of the enzyme replacement therapy. However, beneficial delivery of hLAMAN to the brain requires that the hLAMAN are able to cross the blood-brain-barrier (BBB).

WO 02/099092

PCT/DK02/00388

7

In a preferred embodiment, the present invention involves a treatment method in which a cellular barrier such as the blood-brain-barrier (BBB) is crossed whereby the material is delivered to the target cells.

- 5 Effective enzyme replacement therapy of alpha-mannosidosis patients with recombinant human LAMAN (rhLAMAN) will require the uptake of an active enzyme into the target cells, and furthermore that the rhLAMAN inside the cell is transported to the lysosomes. As described this can be accomplished by tagging the rhLAMAN. The enzyme may e.g. be tagged with mannose. E.g. endothelial cells have a mannose receptor mediated protein
- 10 uptake, the result being that mannose tagged proteins are internalised by liver endothelial cells, but not by cells that do not have this receptor. A number of other tags are conceivable. One particular interesting type of tag is the mannose-6-phosphate (M-6-P). This tag is particular interesting since all cell-types contain this receptor, which is circulating between the Golgi apparatus, the endosomes and the plasma membrane.
- 15 Extracellular enzymes are captured by the M-6-P receptor found on the cell surface and be internalized and delivered to the lysosomes (Willingham, et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 6967). It is established that the secreted forms of lysosomal enzymes produced in mammalian cell lines, for example CHO cells, contain the mannose-6-phosphate tag. This should also be valid for lysosomal alpha-mannosidase, since it is
- 20 normally transported to the lysosomes by a mannose-6-phosphate dependent pathway.

- However, enzyme replacement therapy is not effective against the acute neuronopathic variant of Gaucher disease (Prows et al. Am J Med Genet 71:16-21). To be able to deliver rhLAMAN to the brain a vehicle that can facilitate passage of the blood-brain-barrier (BBB)
- 25 is needed since rhLAMAN is not likely to be able to traverse over the BBB by it self. Such "vehicle" can be a polyamine-modified, a peptide-modified, an antibody-modified LAMAN or even a cell.

DEFINITIONS

30

- By the term "catalyst" is herein meant either the relevant enzyme which is substituted as it is, or an enzymatically equivalent part or analogue thereof. One example of an enzymatically equivalent part of the enzyme could be a domain or sub-sequence of the enzyme which includes the necessary catalytic site to enable the domain or sub-sequence
- 35 to exert substantially the same enzymatic activity as the full-length enzyme. An example of an enzymatically equivalent analogue of the enzyme could be a fusion protein which includes the catalytic site of the enzyme in a functional form, but it can also be a homologous variant of the enzyme derived from another species. Also, completely

WO 02/099092

PCT/DK02/00388

8

synthetic molecules, which mimic the specific enzymatic activity of the relevant enzyme, would also constitute "enzymatic equivalent analogues".

By the term "hybrid molecule" is herein meant a fusion protein between the mhLAMAN
5 catalyst and the peptide required for transport of the vehicle over the BBB and/or cellular membranes. In the said fusion protein the vehicle part can be attached to either the C-terminal or N-terminal end of the mhLAMAN catalyst.

By the term "recombinant techniques" is herein meant a technique which involves
10 recombinant DNA molecules, that is hybrid DNA sequences comprising at least two DNA sequences, the first sequence not normally being found together in nature with the second.

By the terms "recombinant polynucleotide" or "recombinant polypeptide" as used herein means at least a polynucleotide or polypeptide which by virtue of its origin or manipulation
15 is not associated with all or a portion of the polynucleotide or polypeptide with which it is associated in nature and/or is linked to a polynucleotide or polypeptide other than that to which it is linked in nature.

The term "cloning vehicle" describes a plasmid, virus, retrovirus, bacteriophage, cosmid,
20 artificial chromosome (bacterial or yeast), or nucleic acid sequence which is able to replicate in a host cell, characterized by one or a small number of restriction endonuclease recognition sites at which the sequence may be cut in a predetermined fashion, and which may contain an optional marker suitable for use in the identification of transformed cells, e.g., tetracycline resistance or ampicillin resistance. A special class of cloning vectors
25 possess the features necessary for it to operate as an expression vector. An "expression vector" has a promoter positioned upstream of the site at which the sequence is cut for the insertion of the heterologous DNA sequence, the recognition site being selected so that the promoter will be operatively associated with the heterologous DNA sequence. A heterologous DNA sequence is "operatively associated" with the promoter in a cell when
30 RNA polymerase which binds the promoter sequence transcribes the coding sequence into mRNA which is then in turn translated into the protein encoded by the coding sequence.

By the term "target cell" is herein meant a cell or group of cells (tissue) to which the enzymes should be delivered.

35

DETAILED DISCLOSURE OF THE INVENTION

The present invention relates to a cell capable of producing recombinant human LAMAN (rhLAMAN), said cell comprising the 3066 basepair EcoRI - XbaI fragment of a human

WO 02/099092

PCT/DK02/00388

9

cDNA which codes for a human LAMAN protein in which the position corresponding to position 186 of the full length hLAMAN protein is Aspartic acid (Asp, D). In particular, the invention relates to a cell comprising a DNA sequence which codes for the amino acid sequence shown in SEQ ID NO 12.

5

A specific embodiment of the invention relates to a cell obtained by insertion of a DNA fragment comprising the 3066 basepair EcoRI - XbaI fragment of the DNA fragment having the sequence shown in SEQ ID NO 2, in particular a cell obtained by use of the expression plasmid pLamanExp1 having SEQ ID NO. 2. The cell according to the invention may be
10 obtained by transfection of a non-human mammalian cell line such as by transfection of chinese hamster ovary (CHO) cells.

A presently preferred embodiment of the invention relates to a cell obtainable by the culture of the human LAMAN production cell line DSM ACC2549 which has been deposited
15 at the DSMZ for the purposes of patent deposit according to the Budapest Treaty on 6 June 2002. The invention further relates to an expression plasmid pLamanExp1 having the sequence shown in SEQ ID NO. 2.

In another embodiment, the invention relates to a method for the preparation of
20 recombinant human LAMAN, the method comprising a) introducing, into a suitable vector, a nucleic acid fragment comprising a DNA fragment which codes for the amino acid sequence shown SEQ ID NO 12; b) transforming a cell with the vector obtained in step a); c) culturing the transformed host cell under conditions facilitating expression of the nucleic acid sequence; d) recovering the expression product from the culture.

25

The method may further comprise a fermentation step and/or a purification step. In particular, the invention relates to a method wherein the nucleic acid fragment comprises the 3066 basepair EcoRI - XbaI fragment of the DNA fragment shown in SEQ ID
NO 2.

30

An important aspect of the invention relates to a rhLAMAN produced by the method of the invention and to the use of the rhLAMAN according to the invention for the preparation of a medicament for the treatment of alfa mannosidosis.

35 In its broadest aspect, the present invention relates to a recombinant polypeptide and to a nucleotide sequence encoding the LAMAN polypeptide, to an expression system capable of expressing the polypeptide as well as to pharmaceutical compositions comprising the polypeptide or part of it and to the use of the polypeptide for preventing or treating the

WO 02/099092

PCT/DK02/00388

10

development of symptoms related to alpha-mannosidosis(alpha-mannosidosis) caused by a deficiency, in a subject, of the human Lysosomal alpha-mannosidase (hLAMAN) enzyme.

Preferably, the hLAMAN catalyst is administered over cellular membranes, to a target cell,
5 by uptake of hLAMAN into the target cell by taking advantage of a mannose- or M-P-6-
receptor-mediated uptake.

Mannose-6-phosphate tagged hLAMAN is preferably made in a mammalian cell system
(e.g. CHO, COS cells or BHK cells (Stein et al. J Biol Chem.1989, 264, 1252-1259) to
10 secure correct mannose-6-phosphate tagging on the molecule, which ensures efficient
receptor mediated uptake. Mannose-6-phosphate tagged mhLAMAN is secreted into the
medium and purification of rhhLAMAN may be facilitated by the use of ammonium salts
(NH₄Cl) in the fermentation step.

15 Example 3 provides an example of such production of mannose-6-phosphate tagged
hLAMAN

The production of recombinant hLAMAN which is mannose-6-P comprises one or several or
all of the following steps (an outline):

20

A. Synthesis of rhLAMAN

Cloning of specific rhLAMAN

B. Transfection

25 2-10 µg rhLAMAN hybrid vector DNA is used for transfection by phosphate precipitate/
glycerol shock methodology, into mammalian cells (e.g. CHO, COS cells or BHK cells).
Transfection might also be done with an electric shock methodology.

C. Expression of rhLAMAN

30 Synthesis of the active protein and the mannose-6-P modification of said protein is done
during the expression in the mammalian cell system.

D. Purification of rhLAMAN

rhLAMAN is purified using a 6 step procedure - see example 3.

35

E. Test system for mannose-6-P receptor mediated uptake

WO 02/099092

PCT/DK02/00388

11

The ability of produced rhLAMAN to be active in a mannose-6-P receptor mediated uptake is tested by incubating rhLAMAN with normal fibroblasts or fibroblasts from alpha-mannosidosis patients. Uptake into cells is assayed by increased hLAMAN activity.

5 Example 3 shows an example of mannose-receptor-mediated uptake of hLAMAN into cells.

In the following different delivery techniques of LAMAN enzyme across the BBB and/or cellular membranes are described.

10 1) Peptides and proteins as vehicles for passage LAMAN to the target cells by passage over cell membranes and/or the BBB:

A number of studies in animals have shown that certain proteins and/or peptides may act as vehicles for passage of BBB. For instance proteins modified by the insulin fragment (Fukuta et al. Pharmacol Res 11: 1681-1688) or antibodies to the transferrin receptor

15 (Friden et al. Proc Natl Acad Sci USA 88: 4771-4775) can pass the blood-brain barrier. Also proteins modified by coupling to polyamines (Poduslo and Curran. J Neurochem 66: 1599-1606) have been reported to pass the blood-brain barrier.

20 2) Toxins as vehicles for passage LAMAN to the target cells by passage over cell membranes and/or the BBB:

Different bacteria, plants and animals produce toxins. Toxins have many different targets such as the gut (enterotoxins), nerves or synapses (neurotoxins). Toxins can traverse cell membranes via receptor mediated processes and the embodiment of the present invention
25 is to use toxins as vehicles to passage rhLAMAN to the target cells over cellular membranes and/or the BBB. The preferred target cells are cells in the CNS and/or the peripheral nervous system.

A further embodiment of the present invention is that only the peptide pertaining to the
30 translocation over cellular membranes and/or the BBB of the toxin is used.

One example of a toxin used as a vehicle is a bacterial toxin such as Diphtheria Toxin (DT), from the *Corynebacterium Diphtheriae*. Bacterial toxins exhibit a wide range of toxicities and they fall into groups by structure and function. The toxin binds to a target cell and
35 enters the cell via a receptor, and is reduced to separate fragments. The processed toxin can be divided into the following 3 domains: The catalytic domain (C), the receptor domain (R), and the translocation domain (T).

WO 02/099092

PCT/DK02/00388

12

The catalytic fragment and the receptor fragment of the toxin or fragments thereof are replaced by the LAMAN. This fusion protein can traverse cellular membranes and/or the BBB and thereby deliver the LAMAN to the target cells. One example of the engineering of a hybrid molecule could be by recombinant technology

5

A. Synthesis of rhLAMAN

Cloning of specific rhLAMAN cDNA from human liver, placenta or spleen library.

B. Synthesis of DT cDNA

10 1. Cloning of Diphtheria Toxin cDNA.

2. Removal of the cDNA coding for the catalytic domain and the receptor domain or fragments thereof. The remaining fragment of DT is in the following termed "DT(T).

C. Construction of the DT(T)-rhLAMAN hybrid

15 1. Ligation of rhLAMAN cDNA with the DT(T) fragment.

D. Expression

1. Ligate the DT(T)-rhLAMAN construct in a vector for efficient expression in a mammalian cell system e.g. COS, BHK or CHO.

20

2. Conversion of the rLAMAN to mannose-6-P tagged protein occur during the expression in the mammalian cell system.

Other examples of bacterial toxins to be used as vehicles are Clostridium Botulinum,

25 Pseudomonas Exotoxin A produced by *Pseudomonas aeruginosa*, Cholera Toxin produced by *Vibrio cholerae*, and Pertussis Toxin produced by *Bordetella pertussis*.

Further examples of toxins used as vehicles are plant toxins selected from the list of the following plant toxins: cholinesterase inhibitors, protease inhibitors, amylase inhibitors, 30 tannins, cyanogenic glycosides, goitrogens, lectin proteins, and lathyrogens, pyrroizidine alkaloids.

Yet further examples of toxins used as vehicle are toxins from shellfish (saxitoxin) and snakes (alpha-bungarotoxin). The skilled person may add further examples in light of the

35 details and characteristics of the invention.

WO 02/099092

PCT/DK02/00388

13

3) Proteins and/or peptides isolated from bacteria or viruses as vehicles for passage
rhLAMAN to the target cells by passage over cell membranes and/or the BBB:

The transacting element and/or transacting protein from bacteria or viruses can be used
5 together with rhLAMAN to cross the BBB and/or cellular membranes.

Examples of bacteria are selected from the following list: *Neisseria meningitidis*, *S. pneumoniae*,
Hemophilus influenzae, *Staphylococcus* species, *Proteus* species, *Pseudomonas* species, *E.*
coli, *Listeria monocytogenes*, *M. Tuberculosis*, *Neurospora*, *Spirochetes* *Borrelia burgdorferi*
10 from *Iodex ricinus*.

Examples of viruses are selected from the following list of virus families: *Parvoviridae*,
Papovaviridae, *Adenoviridae*, *Herpesviridae*, *Poxviridae*, *Picornaviridae*, *Reoviridae*,
Togaviridae, *Arenaviridae*, *Coronaviridae*, *Retroviridae*, *Bunyaviridae*, *Orthomyxoviridae*,
15 *Paramyxoviridae*, and *Rhabdoviridae*. Relevant examples of viruses selected from the list
mentioned above are e.g. Measles virus, Papova virus, and JC virus.

4) Mannose-receptor mediated uptake of hLAMAN into cells:
Several human cells (monocytes, fibroblasts, lymphocytes) have shown to be able to cross
20 the BBB by it-self (Hickey WF, Kimura H. *Science*. 1988, 239, 290-292, Hickey WF et al. *J*
Neurosci Res. 1991, 28, 254-260). These cells can be "loaded" with rhLAMAN and can
act as a vehicle for transport of rhLAMAN to the brain. Preferably, lymphocytes are used as
vehicles, because they have a long half-life (2-3 months). Uptake of rhLAMAN into
lymphocytes will take advantage of a mannose-receptor-mediated uptake.

25 Mannose-6-phosphate tagged rhLAMAN is made in a mammalian cell system (e.g. CHO,
COS cells or BHK cells (Stein et al. *J Biol Chem*. 1989, 264, 1252-1259) to secure correct
mannose-6-phosphate tagging on the molecule, which ensures efficient receptor mediated
uptake. Mannose-6-phosphate tagged rhLAMAN is secreted into the medium and
30 purification of rhLAMAN may be facilitated by the use of ammonium salts (NH_4Cl) in the
fermentation step.

The disease which is the target for the inventive method is alpha-mannosidosis, and
therefore the catalyst is hLAMAN or an enzymatically equivalent part or analogue thereof.
35 It is most preferred that the catalyst is a recombinant form of the human hLAMAN enzyme
or of the enzymatically equivalent part or analogue thereof, since recombinant production
will allow large-scale production which, with the present means available, does not seem
feasible if the enzyme would have to be purified from a native source.

WO 02/099092

PCT/DK02/00388

14

Furthermore, use of such catalysts, as described herein, for the preparation of pharmaceutical compositions for treatment of the above-discussed diseases is also part of the invention.

Thus, the present invention represents an important advance in the treatment of genetic
5 and/or acquired metabolic brain disorders of alpha-mannosidosis problems associated with prior treatment methods, including the gene therapy and bone marrow transplantation.

The present invention also relates to the new and surprising concept of use as a carrier of a peptide or protein from a structure capable of crossing a biological barrier, such as a
10 cellular barrier including the blood-brain-barrier or a specific membrane of a cell. The object of the carrier function is to deliver an enzyme to a target cell. The target cell is generally a cell wherein the enzyme activity is insufficient either due to an decrease activity of the enzyme or to a situation where an increased activity is desired.

15 In one embodiment, the invention relates to a method for increasing the content of an enzyme in a cell comprising delivery of the enzyme to the tissue relevant for the cell and/or to the cell by use of a protein or peptide capable of crossing a cellular barrier, the protein or peptide being derivable from the group of toxins, bacteria, from viral peptides, and from fragments and modifications thereof.

20

The disease alpha-mannosidosis has been explained in detail above and the present invention is of particular importance in connection with alleviating the progression of symptoms caused by the enzyme defect of hLAMAN. Accordingly, in a preferred embodiment, the invention relates to a method for preventing or treating the development
25 of symptoms related to alpha-mannosidosis comprising the administration of the enzyme recombinant human lysosomal alpha-mannosidase (rhLAMAN) to the tissue relevant for the cells and/or to the cells by use of a protein peptide capable of crossing a cellular barrier or by use of a human cell as a vehicle for delivering of the enzyme to the tissue relevant for the cells and/or to the cells.

30 Similar to the use of a proteins such as insulin or transferrin antibodies as carrier proteins, the use of a protein or peptide derivable from a toxin protein has been suggested as a carrier protein the present invention. Such toxins may be a toxin selected from plant toxins, bacterial toxins and from toxins from animals. The toxin may be modified in order to increase the desired properties of the peptide.

35

In one aspect, a protein or peptide derivable from a virus may be used. However, as virus in its nature has the cell nucleus as target organ for the infection, it is contemplated that a modification or specific fragment not having this effect may be a preferred embodiment.

WO 02/099092

PCT/DK02/00388

15

However, the delivery of an enzyme to an enzyme deficient cell related to alpha-mannosidosis by use of an protein virus is a further surprising aspect of the invention.

As virus in general is an inhomogeneous group with respect to affinity for different cells, it
5 is within the spirit of the present invention to select the concrete virus in accordance with the desired target cell to which the enzyme is to be delivered.

Affection of the nervous system, especially of the central nervous system, very often plays a significant role in the broad range of diseases caused by enzyme deficiency. Accordingly,
10 in a preferred embodiment, the carrier protein or peptide is one which is capable of crossing the blood-brain-barrier.

This is possible by use of proteins or peptide which due to their nature bears a structure relevant for such transduction. In an important method according to the invention, the
15 cellular barrier includes the blood-brain-barrier (BBB) and the tissue is a tissue and/or cell of the central nervous system.

The overall idea of the present invention is to prepare a construct comprising the enzyme in question the delivering protein/peptide. Accordingly, in one aspect the invention relates
20 to a method wherein the enzyme forms a hybrid with the protein or peptide capable of crossing the cellular barrier.

As explained in further details in the text of the present specification the hybrid is preferably produced by recombinant techniques. However, the construct may be produced
25 by techniques of protein synthesis generally known in the art including solid phase synthesis. The complete hybrid or part of the hybrid may accordingly be produced synthetically or a part of the hybrid construct may be produced by use of a genuine protein or peptide. The enzyme part and delivery part may be linked by different techniques known in the art.

30

Preferably, the hLAMAN is made by recombinant techniques. In a further embodiment, the LAMAN is human (hLAMAN) and still more preferred mature human hLAMAN (mhLAMAN) or a fragment thereof. The fragment may be modified, however the active sites of the enzyme should be preserved.

35

In a still further aspect of the invention, cells are used as a vehicle for delivering an enzyme, preferably hLAMAN to the target cell. The preferred human cell is selected from human monocytes, human fibroblasts, and human lymphocytes. As explained above, it is preferred for target cells of the central nervous system that the cell for delivering the

WO 02/099092

PCT/DK02/00388

16

enzyme is capable of crossing the BBB for delivering the hLAMAN to the tissue and/or cells of the central nervous system.

- In order to facilitate the delivery of the enzyme to the target cell, the hLAMAN may be
- 5 transferred to a target cell by means of a mannose-receptor-mediated uptake. Such uptake may be further increased when the hLAMAN is a mannose-6-P tagged hLAMAN, preferable made by expressing hLAMAN in a mammalian cell system. Preferred mammalian cell system is selected from CHO cells, COS cells, and BHK cells.
- 10 In general, the target cell is a cell wherein the activity, such as hLAMAN is insufficient for the optimal function of the cell. Insufficient activity of hLAMAN may be measured by one or more of the parameters selected from monitoring increased levels of mannose-rich oligosaccharides in urine, analysis of hLAMAN activity in material from the patient such as in leukocytes and/or in skin fibroblasts, presence of clinical symptoms or increase in rate of
- 15 development of clinical symptoms of alpha-mannosidosis.

- A significant feature of insufficient hLAMAN activity is a cell wherein a massive intracellular accumulation of mannose-rich oligosaccharides is present. Naturally, such cell is a target cell according to the present invention. The target cell may also be cells of the nervous
- 20 system. In addition, target cells for delivering the enzyme also includes one or more cell types selected from human monocytes, human fibroblasts, human lymphocytes, human macrophages.

- An increased activity of the hLAMAN may be used as a parameter for a treatment schedule
- 25 an may be measured by one or more of the parameters selected from monitoring increased levels of mannose-rich oligosaccharides in urine, analysis of hLAMAN activity in material from the patient such as in leukocytes and/or in skin fibroblasts, presence of clinical symptoms or increase in rate of development of clinical symptoms of alpha-mannosidosis.

- 30 It is a very important aspect of the invention to perform the treatment of a possible enzyme defect prenatally. In a further aspect of the invention, the cellular membrane is the fetal-maternal barrier (placenta). It is also within the scope of the invention to deliver the enzyme-protein-construct directly to the fetus prenatally.
- 35 In a still further embodiment, the invention relates to an antibody raised against any of the constructs formed by any of the enzymes and any of the proteins and/or peptide mentioned herein. Such antibody may be used for the targeting of the construct, for inactivation of the construction including increasing the elimination of the construct from the subject or even more important for detecting or monitoring the level of intracellular

WO 02/099092

PCT/DK02/00388

17

levels of LAMAN. The antibody may be a polyclonal antibody or a monoclonal antibody and by be produced by techniques known in the art.

Any enzyme construct disclosed above as well as the antibody thereto may be use for the
5 preparation of a medicament for a treatment in accordance with the method disclosed in the present context. Accordingly, the present invention also relates to a pharmaceutical medicament comprising an enzyme linked to a carrier system such as a protein or peptide or to a cell system as disclosed in detail above.

10 Accordingly, in one aspect, the present invention relates to a construct as well as to the use of a construct comprising an enzyme and/or a protein or peptide protein peptide capable of crossing a cellular barrier or a human cell as a vehicle for delivering of the enzyme to the tissue relevant for the cells and/or to the cells as disclosed in any of the methods mentioned and specified in the claims for the preparation of a medicament.

15

Also the antibody and use thereof the preparation of a medicament is within the scope of the invention.

In a presently preferred aspect, the present invention relates to the use of the LAMAN
20 coding sequence of figure 4 for the use in gene therapy. It is suggested that a recombinant viral or non-viral vector comprising the cDNA coding for the biologically active human LAMAN and able to infect and/or transfect and sustain expression of the biologically active human LAMAN in mammalian cells could be produced. Such constructs could be used to treat alpha-mannosidase patients.

25

In a very one important aspect the present invention the principle of "gene therapy" is developed in relation to therapy by bone marrow transplantation. Bone marrow transplantation (BMT) with cells that do produce LAMAN is known to have a curative effect on individuals (test animals) suffering from alpha-mannosidosis (Walkley et al. Proc. Nat.
30 Acad. Sci. 91: 2970-2974, 1994). BMT may even be beneficial to human patients. In 1987 a child with mannosidosis was treated with BMT (Will et al. Arch Dis Child 1987 Oct;62(10):1044-9). He died after 18 weeks due to procedure related complications. This disappointing result could be explained by heavy immunosuppressive treatment before death. This treatment was necessary because the patient received non-self bone-marrow
35 cells. One way to overcome the problem of rejection of foreign bone marrow cells could be to use the patients own cells. We suggest to transfect such stem cells with a construct comprising the LAMAN cDNA nucleic acid sequence to obtain bone marrow stem cells that express enhanced levels of LAMAN, and finally to reintroduce these cells into the bone marrow of the patient.

WO 02/099092

PCT/DK02/00388

18

5

LEGEND TO FIGURES

Figure 1.

10 Description of oligonucleotides used for PCR amplification.

The two oligonucleotides ICO1106 and ICO1107 are shown below the LAMAN sequence.

Numbering refers to the numbering in the GenBank sequence Accession # U60266.1. The highlighted regions in the LAMAN sequence represent the regions of homology with the oligonucleotide primers. The asterisk denotes the C to A change introduced at position 53

15 to generate a *Xho* I site for ease of manipulation. This is a silent mutation in the codon for the arginine residue at position 13 in the amino acid sequence of the α -mannosidase protein. The triplets coding for this arginine residue and for the termination codon are boxed. The PCR primer ICO1107 has a 5' extension incorporating a *Xba* I site to allow cloning of the amplified DNA.

20

Figure 2.

Strategy for PCR cloning of LAMAN.

Figure 3.

25 Map of the expression plasmid pLamanExp1.

"pCMV" is the Human cytomegalovirus immediate-early promoter/enhancer. "Intron" is a chimeric Intron. "rhLAMAN" is the recombinant human lysosomal alpha monnosidase coding region. "SV40-term" is the SV40 late termination and polyadenylation signal. "F1" is Phage f1 region. "SV40 Eprom" is the SV40 early promoter/enhancer and origin. "DHFR"

30 Is the Mouse dihydrofolate reductase coding region. "pA" is a synthetic polyadenylation signal. "AmpR" is the beta-lactamase (AmpR) coding region.

Figure 4.

Map of the expression plasmid pAsaExp1.

35 "pCMV" is the Human cytomegalovirus immediate-early promoter/enhancer. "Intron" is a chimeric Intron. "rhASA" is the recombinant human Arylsulfatase A coding region coding region. "SV40-term" is the SV40 late termination and polyadenylation signal. "F1" is Phage f1 region. "SV40 Eprom" is the SV40 early promoter/enhancer and origin. "DHFR" is

WO 02/099092

PCT/DK02/00388

19

the Mouse dihydrofolate reductase coding region. "pA" is a synthetic polyadenylation signal. "AmpR" is the beta-lactamase (AmpR) coding region.

Figure 5. SDS-PAGE analysis of anion exchange chromatography for rhLAMAN. See table 2
5 for content of each lane.

Figure 6. SDS-PAGE analysis of anion exchange chromatography for rhLAMAN (larger
scale). See table 3 for content of each lane.

10 Figure 7. Fibroblasts from a patient with mannosidosis (GM00654) loaded with DEAE-
purified rhLAMAN for 40 hours in complete medium.

Figure 8. Fibroblasts from a patient with mannosidosis (GM00654) loaded with DEAE-
purified rhLAMAN for 40 hours in complete medium with the addition of 5 mM Mannose-6-
15 Phosphate (M-6-P) or 5 mM Glucose-6-Phosphate (G-6-P).

Figure 9. Fibroblasts from a patient with alpha-mannosidosis (GM00654) loaded with
DEAE-purified rhLAMAN for 40 hours in complete medium with the addition of 5 mM
Mannose-6-Phosphate (M-6-P) or 5 mM Glucose-6-Phosphate (G-6-P). Endogenous LAMAN
20 activities are subtracted from each bar.

Figure 10. Fibroblasts from a patient with alpha-mannosidosis (GM00654) loaded with
DEAE-purified rhLAMAN for 40 hours in complete medium with the addition of 5 mM
Mannose-6-Phosphate (M-6-P) or 5 mM Glucose-6-Phosphate (G-6-P).

25

Figure 11. Fibroblasts from a patient with alpha-mannosidosis (GM00654) loaded with
DEAE-purified rhLAMAN for 40 hours in complete medium with the addition of 5 mM
Mannose-6-Phosphate (M-6-P) or 5 mM Glucose-6-Phosphate (G-6-P) with the endogenous
LAMAN activities subtracted from each bar.

30

EXAMPLES

Example 1: Cloning of hLAMAN cDNA and construction of expression vector

35 PCR and cloning

The source of the PCR DNA template was a HepG2 cDNA library made from mRNA isolated
from the HepG2 cell line (ATCC # HB-8065) using the Superscript Plasmid System
manufactured by Gibco BRL, catalogue # 18248-013. Two hundred ng of DNA from this
library was amplified using KlenTaq polymerase mix (Clontech, catalogue # 84171-1) in a

WO 02/099092

PCT/DK02/00388

20

50 ul reaction volume containing 0.2mM dNTPs and 0.4 uM each of ICO1106 (5' TCTGCGC
TCGAGGCTGCCTGGACTCA, SEQ ID NO. 3) and ICO1107 (5' TGCTAATCTAGAGGG
CCCATCCCAGCAGAC, SEQ ID NO. 4). Two-cycle PCR was used with an initial denaturation
step at 96°C for 2 minutes followed by 30 cycles of 94°C for 30 seconds and 65°C for 4.5
5 minutes. A final extension of 10 minutes at 72°C was used at the end to ensure that the
extension products were filled out.

The PCR reaction mix was run out on a gel and a PCR product of 3038 bp was isolated.
One twentieth of this material was re-amplified with the same two primers as described
10 above, except that the temperature of the extension reaction was 68°C. The 3038 bp
amplified fragment was gel isolated, digested with *Xho* I and *Xba* I and assembled into
plasmid pAsaExp1 (Fig 5 and 6) cut with *Eco*R I and *Xba* I together with oligos ICO1108
(5'AATTCGCCGCCATGGGCGCCTACGCGCGGGCTTCGGGGGTCTGCGC) and ICO1109
(5'TCGAGCGCAGACCCCGAAGCCCG CGCGTAGGCGCCCATGGCGGCG) in a three part
15 ligation (Fig.2).

Sequencing

Seven plasmid clones from the ligations described above were sequenced with the Big dye
terminator cycle sequencing kit from PE/ABI (catalogue # 4314417). Two vector primers,
20 ICO873 (5' TGCAGCTTATAATGGTTACA, SEQ ID NO. 5) and ICO929 (5'
GGCACCTATTGGTCTTACTG, SEQ ID NO. 6) and five LAMAN-specific primers, ICO1110 (5'
AAGTCGTGCGAGACCTTGTG, SEQ ID NO. 7), ICO1111 (5' CCGCCAAGGA ATCTGTGCTG,
SEQ ID NO. 8), ICO1112 (5' TACGAGCGCCTCAGCTACAA, SEQ ID NO. 9) and ICO1113 (5'
GCGGAAGGTGAATTGGATGG, SEQ ID NO. 10) and ICO1114 (5' GCCTCAGGTGCC
25 TACATCTT, SEQ ID NO. 11) were used. The results are summarized in Table 1.

Construction of the correct, full-length LAMAN cDNA

As seen from Table 1, none of the seven cDNA clones match the published sequence
(Nilssen et al. Hum.Mol.Genet. 6, 717-726 1997, revised version Accession # U60266.1).
30 In addition to a number of random changes they all have the TA to AT changes at positions
1670 and 1671 in the plasmid sequence (SEQ ID NO:1). This results in a valine to
asparatic acid substitution in residue 186 of the LAMAN protein sequence, see SEQ ID NO.
12. The same valine to asparatic acid change is also reported by two other research
groups (Liao et al. J.Biol.Chem. 271, 28348-28358, 1996 and Nebes et al. Biochem
35 Biophys. Res. Commun. 200, 239-245, 1994). Hence, for the purpose of this study, a full-
length LAMAN sequence was patched together from plasmid clones pLaman-50 and
pLaman-70 using the unique SanD I site at position 2388 (Figs. 2 and SEQ ID NO:2).

A 1287 bp *Eco*R I – SanD I fragment from pLaman-50 and a 1779 bp

WO 02/099092

PCT/DK02/00388

21

SanD I – Xba I fragment from pLaman-70 were isolated and assembled together in plasmid pAsaExp1 (Fig. 5 and 6) cut with EcoR I and Xba I in a three part ligation (Fig. 2). The resulting plasmid is called pLamanExp1. The LAMAN insert in this plasmid was confirmed by sequencing and found to match the sequence (Nilssen et al. Hum.Mol.Genet. 5 6, 717-726, 1997 accession # U60266.1) except at amino acid position 186 as referred to in the plasmid sequence shown in SEQ ID NO:2. The map of this plasmid is shown in Fig. 3.

WO 02/099092

PCT/DK02/00388

22

Table 1. Summary table of sequencing results

Clone #	Nucleotide change	Amino acid change
27	1287, G to A	58, V silent
	1292, C to T	60, P to L
	1320, T to G	69, P silent
	1670, T to A	186, V to D
	1671, A to T	186, V to D
	2048, G to T	682, L to I
	2829, C to T	572, A silent
	3423, C to T	770, P silent
	3448, G to A	779, N to D
	3826, C to T	905, L silent
	3835, T to C	908, W to R
	4112, T to C	1000, F to S
46	1258, G to A	49, A to T
	1670, T to A	186, V to D
	1671, A to T	186, V to D
	2360, A to G	416, E to G
	2415, A to G	434, A silent
	3409, A to G	766, N to D
50	1670, T to A	186, V to D
	1671, A to T	186, V to D
	2401, T to C	430, S to P
	2465, G to A	451, G to D
	2824, C to T	571, P to S
	2886, C to T	591, R silent
	3075, C to T	654, D silent
	3281, A to G	723, D to G
	3423, C to T	770, P silent
	3488, C to T	792, T to I
66	1165, T to A	18, S to T
	1670, T to A	186, V to D
	1671, A to T	186, V to D
	2450, A to G	446, H to R
	2686, A to G	525, M to V

WO 02/099092

PCT/DK02/00388

23

	3368, T to C 3423, C to T	752, I to T 770, P silent
70	1391, A to G 1670, T to A 1671, A to T 1869, C to T 1976, A to G 2048, C to T 2146, A to G 3186, G to A 3423, C to T	93, Q to R 186, V to D 186, V to D 252, L silent 288, N to S 312, I to L 345, K to E 691, Q silent 770, P silent
74	1670, T to A 1671, A to T 1823, A to G 1849, A to G 1910, A to G 2012, C to T 2048, C to T 2284, T to G 3330, A to G 3558, A to T 3801, G to T 3813, G to A 3828, G to A	186, V to D 186, V to D 237, Q to R 246, K to E 266, N to S 300, A to V 312, I to L 391, Y to D 739, T silent 815, R silent 896, L silent 900, V silent 905, L silent
76	1169, C to T 1670, T to A 1671, A to T 2653, G to A 3008, A to G 3724, C to T	19, A to V 186, V to D 186, V to D 514, V to D 632, Q to R 871, L silent

NOTE: The clones are in plasmid pLamanExp1. The numbering is as shown in the plasmid sequence (SEQ ID NO:2). Identical changes between clones are highlighted in different
5 colors. The reference sequence for the lysosomal alpha-mannosidase cDNA has the GenBank Accession number U60266.1 (1, revised).

WO 02/099092

PCT/DK02/00388

Discussion

The inserts in all seven of the clones were confirmed to be LAMAN by sequence analysis.

As seen in Table 1 a number of changes were found in all of the clones when compared to

5 our reference sequence (1, revised version accession # U60266.1). Most of them are likely to be random PCR errors. Curiously, the TA to AT change at positions 1670 and 1671 resulting in a valine to asparatic acid substitution was common to all the clones. The two other sequences published for LAMAN (Nebs et al. Biochem. Biophys. Res. Commun. 200, 239-245, 1994, Liao et al. J.Biol.Chem. 271, 28348-28358, 1996) also report asparatic
10 acid in that position. In the light of these observations a full-length correct sequence was put together with the help of the SanD I site at position 2388 as described under materials and methods. The sequence of the cDNA insert in this clone is identical to the reference sequence except for the valine to asparatic acid change mentioned above.

15 Evaluation and conclusions

A combination of PCR amplification and Incorporation of synthetic oligonucleotides assembled a full-length clone of LAMAN. It matches the sequence Accession # U2066.1, our reference sequence except in the valine to asparatic acid substitution alluded to earlier in this report. This LAMAN cDNA sequence also carries a C to A change at nucleotide
20 position 1152 (see plasmid sequence in SEQ ID NO. 2). This was engineered to create a Xho I site for ease of manipulation and does not change the primary sequence of the LAMAN protein. It also contains a G to A change at position 3186 and a C to T change at position 3423 with no effect on the primary amino acid sequence of the LAMAN protein.

25 All seven of our clones had the valine to asparatic acid substitution in the 186th residue of the protein. Since the same substitution was also reported by two other groups (accession #s U68567 and U05572, refs. 2 and 3) the asparatic acid residue was left unchanged at that position. The reference sequence Accession # U60266.1 differs from the two other published sequences mentioned above at a few other positions. When compared to
30 sequence Accession # U68567 (Liao et al. J.Biol.Chem. 271, 28348-28358, 1996) three other changes can be found. Curiously, the threonine to isoleucine change occurring at amino acid position 312 (Liao et al. J.Biol.Chem. 271, 28348-28358, 1996) was present in three of our clones (Table 1). None of our clones had either the 3 base-pairs deletion encoding the glutamine residue at position 342 or the G to T silent mutation in the leucine
35 residue at position 753 found in (3, Accession # U05572). Three of our clones however, had a C to T silent change in the proline at position 770.

WO 02/099092

PCT/DK02/00388

25

Example 2: Transfection, selection of producer cell and preliminary characterisation of recombinant human LAMAN

5 Materials and methods:

Materials used for cell culture

Media:

<u>Material</u>	<u>Company</u>	<u>Catalog #</u>
Penicillin/Streptomycin	BioWhittaker	17-602E
10 L-Glutamine, 200 mM	BioWhittaker	17-605E
HT Supplement (100X)	Gibco BRL	11067-030
Ex-Cell 302 Serum-Free Media	JRH BioSciences	14312-78P

Media for DG44.42 Cells:

15 Ex-Cell 302 Serum-Free Medium	500 ml
Penicillin/Streptomycin	5 ml
L-Glutamine (4 mM final concentration)	10 ml
100X HT Supplement	5 ml

20 *Media for ASA Masterwell Cells:*

Ex-Cell 302 Serum-Free Medium	500 ml
Penicillin/Streptomycin	5 ml
L-Glutamine (4 mM final concentration)	10 ml
20 nM Methotrexate	0.5 ml (20 µM MTX*)

25

*Diluted from 20 mM Stock.

Preparation of 20 mM Methotrexate Stock:

30 Materials:

1. Methotrexate, U.S.P. : ICN catalogue # 102299
2. WFI (Water for Injection)
3. 1.0N NaOH to adjust pH.
4. 0.2µm filter unit of appropriate size

WO 02/099092

PCT/DK02/00388

26

Gel electrophoresis:

Gel electrophoresis was done using the Novex system with Nupage Bis/Tris gels run at 200 volts for 1.2 hours. Staining was done with Coomassie brilliant blue R-250 according to the
5 manufacturer's specifications. Total protein was measured by the Bradford method using the BioRad reagent (catalogue # 500-006) and bovine serum albumin as standard.

*LAMAN enzyme assay**Materials:*

- 10 4X assay buffer consisting of:
250 μ M p-Nitrophenyl-alpha-D-mannopyranoside
100 μ g/mL BSA
200 mM Na Acetate (pH 4.5)

15 1.8 M Na_2CO_3

Procedure:

- For screening purposes the assays were done in flat-bottomed ELISA plates. 50 μ l of the 4X assay buffer was added to 150 μ l of sample or an appropriate dilution of it (in 10 mM
20 Tris pH 7.4 containing 150 mM NaCl). The plates were incubated overnight at room temperature, stopped with 100 μ l of 1.8 M Na_2CO_3 and the absorbance recorded at 405 nm on a plate reader.

- For determination of specific activity of the DEAE-purified samples, the assays were set up
25 in tubes with all the volumes doubled. Incubations were at 37°C for periods ranging from 3-60 minutes using 10-10,000 ng of enzyme. The samples were read on a spectrophotometer using a cuvette of 1 cm path length. Specific activity is defined as μ moles of p-Nitrophenyl-alpha-D-mannopyranoside hydrolysed per minute per mg protein.

30

Introduction

To be able to produce rhLAMAN the expression plasmid (pLamanExp1, Fig 3, SEQ ID NO. 2) was cell type the following experiment was performed.

- To obtain the final clonal cell line expressing rhLAMAN, the expression plasmid
35 pLamanExp1 (Fig 3, SEQ ID NO. 2) was transfected into the host cell line DG44.42 (obtained from DG44 by limiting dilution). High LAMAN producing masterwells, as determined by LAMAN activity, were selected.

WO 02/099092

PCT/DK02/00388

27

To further increase the yield, the cells from high LAMAN producing masterwells were subjected to amplification with methotrexate (MTX) as described by Gasser et al. Proc. Natl. Acad. Sci. U.S.A 79, 6522-6526. 1982. After amplification, the best producers were cloned twice by limiting dilution.

5

Transfection conditions and selection of high producing masterwells

1x10⁷ CHO DG44.42 cells were resuspended in 800 µl Hepes Buffered Electroporation Media with 1020 µg pLamanExp1 DNA (linearized with *Ahd I*) +120 µg salmon sperm DNA in a 0.4 cm electroporation cuvette. The cells were electroporated using a Bio-Rad Gene
10 Pulser II set at 250 volts with 500 µF capacitance. The contents of the cuvette were then transferred to a T-75 flask containing 20ml of JRH Ex-Cell 302 Serum-Free Medium + HT (hypoxanthine and thymidine) + 4 mM L-glutamine + 100 units/ml penicillin + 100 µg/ml streptomycin and then placed in a 37 °C, 5% CO₂ incubator. 24 hours later the cells were counted, centrifuged and transferred to two T-flasks, one with 20 nM and the other with 50
15 nM MTX replacing the HT in the growth media described above. A media sample was saved and run in a LAMAN activity assay - see materials and methods. The culture showed activity and was plated in 96-well dishes as follows; 2 plates each at 500 and 1000 cells/well in 100 nM MTX and 5 plates each at 50, 100 and 200 cells/well in 20 and 50 nM MTX respectively. The wells were fed fresh media every 5-7 days. Four weeks post
20 electroporation a LAMAN activity assay was done on media from wells containing 1 to 4 clones. The masterwells with the highest LAMAN activity were then expanded and run in a 6-day LAMAN activity assay. In a 6-day LAMAN assay cells were seeded at 1x10⁵ cells/ml in 5 ml media with MTX in a T-12.5 flask. Six days later the cells were counted. The cells were then centrifuged and the supernatant analyzed in the LAMAN activity assay. Results
25 were normalized for cell number. The 12 best masterwells were then amplified.

Amplification

Six masterwells were amplified starting from 20 nM MTX and six starting from 50 nM MTX. Amplification was done by seeding cells at 1x10⁵ cells/ml in T-25 Falcon flasks, in 10ml of
30 media. The cells were split every 4-6 days and analyzed in a 6-day assay. The cells were then amplified to the next level (50, 100, 200 nM MTX) on the next split as well as the cells being maintained at previous MTX levels. No amplification occurred at 100 and 200 nM MTX. The highest producing masterwell at 50 nM MTX (20L-31) was selected for further subcloning.

35

1st Round Limiting Dilution Cloning

20L-31 was spinner adapted before cloning. The cells were seeded at 1.25, 2.5, 5 and 10 cells/well in 100 µl media containing 10% conditioned media in a flat-bottom 96-well plate. They were fed with 100 µl fresh media every 5-7 days. The wells were scanned for single

WO 02/099092

PCT/DK02/00388

28

clones within 4 weeks and these cells were transferred to 24-well plates with 1 ml fresh media. Growing clones they were transferred to a T-25 flasks, expanded and run in the 6-day LAMAN assay. Twenty-seven clones were analyzed in this fashion. One clone was chosen on the basis of cell growth, viability and activity. The clone chosen, 20L-31.62, had
5 twice the LAMAN activity as the corresponding masterwell.

2nd Round Limiting Dilution Cloning

The first round clone 20L-31.62 was already spinner adapted and was taken through the same subsequent steps of limiting dilution cloning as described above. Twenty-seven
10 clones were analyzed in a 6-day LAMAN assay and one clone was retained as the final clone on the basis of cell growth, viability and LAMAN activity. The clone chosen, 20L-31.62.5, was also twice as active as the masterwell. It was maintained in media containing 50 nM MTX. This clone has been designated "CHO-LAMAN 20L-31.62.5
020118".

15

Characterization of rhLAMAN

The characterization of rhLAMAN for specific LAMAN activity was done with partially purified material from the first round clones.

20

Purification

A preliminary purification of rhLAMAN was attempted for the purposes of determining specific activity and characterise the product by gelelectrophoresis.

25 Briefly, 800 ml of conditioned media from the masterwell, 20L-31 (cultured for in T-flasks for 10 days) was dialyzed against 20 mM Tris, pH 7.6 and loaded on a 5 ml DEAE-Sephacrose column (AP-Biotech, Catalogue # 17-0709-01), equilibrated in the same buffer. The column was washed thoroughly with equilibration buffer containing 20 mM NaCl and then eluted in steps using 120 mM, 250 mM and 1 M NaCl. One ml fractions were collected
30 and analyzed by Bradford protein assay and SDS-PAGE. The gel profile is shown in Fig. 5. The fractions are labeled serially according to the salt concentration used for elution. Two pools were collected. The content of each lane is described in table 2 below.

Table 2: Contents of 4-12% BisTris w MES/SDS reducing gels

Lane	Sample	µg/ml by Bradford	total µg in lane		Lane	Sample	µg/ml by Bradford	total µg in lane
1	Marker				13	250.5	460	9.2
2	Start	124	2.48		14	marker		
3	Flow Through	80	1.6		15	1.1	1023	20.46
4	120.6	1827	36.54		16	1.2	ND	ND
5	120.7	1755	35.1		17	1.3	ND	ND
6	120.8	1250	25		18	1.4	ND	ND
7	120.9	963	19.26		19	1.5	ND	ND
8	120.10	619	12.38		20	1.6	1471	29.42
9	250.1	477	9.54		21	1.7	ND	ND
10	250.2	361	7.22		22	1.8	ND	ND
11	250.3	303	6.06		23	1.9	ND	ND
12	250.4	334	6.68		24	1.10	ND	ND

5

The samples correspond to the fractions collected. The fractions are numbered serially according to the salt concentration in the step gradient used (120 mM, 250 mM, 1 M). ND stands for not done.

The 120 mM pool (lanes 4-8 in Fig. 5) had a specific activity of 0.8 U/mg and 1.25 mg/ml total protein. The 250 mM pool (lanes 9-12 in Fig. 5) had a specific activity of 1 U/mg and 0.48 mg/ml total protein.

10 Another purification was attempted starting with 1.2 L of dialyzed medium from 20L-31 grown in T-flasks applied to a 50 ml DEAE-Sepharose column. After washing the column in the same manner, rhLAMAN was eluted with a steep linear gradient of 20-120 mM NaCl. 5 ml fractions were collected and analyzed by SDS-PAGE. Figure 6 shows the gel profile and the pooled fractions are marked. The content of each lane is described in table 3 below.

Table 3: Contents of 4-12% BisTris w MES/SDS reducing gel

Lane	Fraction	mM NaCl
1	10 µg of rhLaman from purification shown in Fig. 5	120 mM by Step elution
2	Markers	
3	5	40
4	6	40
5	10	50
6	11	50
7	12	50
8	16	60
9	18	70
10	60	140
11	62	140
12	63	150

5 The pooled material (25 ml) had a specific activity of 2 U/mg at a protein concentration of 1 mg/ml.

Conclusion

10 A LAMAN production cell line (CHO DG44.42 subclone 20L-31.62.5) was obtained after MTX amplification and two rounds of cloning. This clone has been designated "CHO-LAMAN 20L-31.62.5 020118" and deposited at the DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, 38124 Braunschweig, 15 GERMANY) for the purposes of patent deposit according to the Budapest Treaty on 6 June, 2002.

The CHO cell-based expression system described here produces in excess of 20 mg/L rhLAMAN, which is secreted into the culture medium as a single-chain molecule. A 20 preliminary purification was attempted. The purification resulted in a partially purified rhLAMAN preparation, see elution profile Fig 5 and 6 that can be used for cellular uptake studies and further purification.

WO 02/099092

PCT/DK02/00388

31

Example 3: In vitro characterisation of rhASA

Materials and methods:

5 Complete medium

MEM Eagle-Earles BSS with

15% FCS

2 mM Glutamine

2X conc. Vitamins

10 2X conc. non-essential AA

2X conc. MEM-AA

1 mM Na-pyruvate

TBS composition

15 10 mM Tris-HCl

150 mM NaCl

Adjust pH to 7.5

Alfa-Mannosidase activity assay protocol

20

*4x assay buffer*16 mM para-Nitrophenyl- α -D-mannopyranoside

2 mg/ml BSA

in 0.4 M NaAc pH 4.5

25

Stop buffer

1.8 M Na₂CO₃

For measuring activity in cell supernatants, the assay was run in microtiter plate. For cell

30 lysates, the assay was performed in Eppendorf tubes, see below.

Microtiter plate set-up

150 μ l of sample at appropriate dilution (in TBS)50 μ l of 4x assay buffer

WO 02/099092

PCT/DK02/00388

32

The plate is incubated at 37°C for 30 minutes.

Stop by adding 100 µl stop buffer and read the plate at OD 405.

Eppendorf tube set-up

5 300 µl of sample at appropriate dilution (in TBS)

100 µl of 4 x assay buffer

The tubes are incubated at 37°C for 30 minutes to 2 hours.

Stop by adding 200 µl stop buffer, centrifuge the tubes (13 200 rpm, 10 minutes) and transfer the supernatant to a cuvette.

10 Measure at OD 405.

The LAMAN activity is then calculated using the millimolar extinction coefficient for the product in the formula:

15 $(OD\ 405 \times V)/(18.8 \times t \times v) = U/ml$

V = total volume for assay (ml)

t = incubation time (min)

v = sample volume (ml)

20 The millimolar extinction coefficient for para-nitrophenol = $18.8\ M^{-1}\ cm^{-1}$

1 U of LAMAN activity is defined as the amount of enzyme that liberates 1 µmol para-nitrophenol/min, 37°C, pH 4.5.

25 BCA protein determination kit

Protein concentration was determined using the BCA Protein assay kit (no. 23225) from Pierce. This kit utilizes the principle of the reduction of Cu^{2+} to Cu^{+} by protein in an alkaline medium (the Biuret reaction). The Cu^{+} ions are then reacted with a reagent containing bicinchoninic acid resulting in a highly sensitive and selective colorimetric detection. The

30 kit was performed according to the manual.

Microtiter plate set-up

10 µl of the BCA-Standards for a standard curve

35 10 µl of BSA reference in duplicate

10 µl of MilliQ water used as blank in at least triplicate

10 µl of sample or diluted sample in MilliQ water

WO 02/099092

PCT/DK02/00388

33

200 µl of BCA assay buffer in all wells

Mix ~30 seconds

Incubate samples for 30 minutes at 37° C

Measure the absorbance within 10 minutes at 562 nm and determine the concentrations

5 from the standard curve.

Introduction

At an early stage, before the purification procedure was developed, the rhLAMAN was characterized in an *in vitro* cell system. For this purpose, LAMAN-containing supernatants from LAMAN-CHO cells were purified using a DEAE purification step. Briefly, LAMAN-containing supernatants from T flasks with the DG44.42 subclone 20L-31.62.5 were diafiltered and concentrated and then added to the column after equilibration with 10 mM Tris-HCl, pH 7.6. After loading, the column was washed with the same buffer and then with 10 mM Tris-HCl + 0.02 M NaCl. 3 column volumes were used for each washing step. The bound proteins were eluted using a NaCl gradient ranging from 0.02 to 0.3 M NaCl. 5 ml fractions were collected. The fraction with the highest specific activity was purified approximately 100 times and the specific LAMAN activity was around 4 U/mg. This material was used in the cell systems described below.

20

1. Cellular uptake of rhLAMAN by MANN fibroblasts

Aim of the study

To get *in vitro* proof of concept regarding cellular uptake of rhLAMAN in fibroblasts from individuals affected by alpha-mannosidosis (MANN fibroblasts).

Experimental design

Fibroblasts from a patient with mannosidosis (MANN fibroblasts GM00654, purchased from Coriell Cell Repository, USA) were grown confluent in 6-well plates in complete medium. Cells were washed with PBS and rhLAMAN (diafiltrated, concentrated and DEAE-purified, see purification part for details) was added (0-200 mU/ml) ± 5 mM Mannose-6-Phosphate (M-6-P) or Glucose-6-Phosphate (G-6-P, control) in complete medium. The specific LAMAN activity of the added enzyme was 4 U/mg total protein. The cells were left for 40 hours and then harvested; medium was removed, centrifuged and buffer was changed to TBS using PD10 columns. Cells were washed with PBS and trypsinated, washed with TBS and then lysed with 0.5 % Triton X-100 in TBS. The lysed cells were centrifuged (13 200 rpm, 10 minutes) and lysate was collected. LAMAN activity and protein concentration were measured in all samples.

WO 02/099092

PCT/DK02/00388

34

For details on the LAMAN activity assay and complete medium composition, see Materials and methods. Protein determination was performed in micro-titer plate according to our internal SOP. See Materials and methods for a brief description of the method.

5

Results and discussion

The MANN fibroblasts had a background LAMAN activity of 1-1.5 mU/mg total protein. DEAE-purified rhLAMAN gave a maximal cellular load of 37 mU/mg total protein (after addition of approximately 75 mU/ml for 40 hours), see Fig. 7. Addition of M-6-P inhibited 10 90 % of the cellular uptake and there was no inhibition using G-6-P. This suggests that the enzyme is taken up mainly by the M-6-P receptor. Results are shown in Fig. 8 and 9.

Example 4: Small-scale cultivation of producer cell

Cell line and culture medium

15 A DHFR minus Chinese Hamster Ovary (CHO) cell line, DG44.42 subclone 20L-31.62.5, expressing rhLAMAN, is routinely cultivated in modified EX-CELL 302 CHO serum-free medium without phenol red (JHR Biosciences Europe, UK). The medium contains 1.6 g/L sodium bicarbonate, 4 mM HEPES and 0.1 % Pluronic®F-68, and is supplemented with 4 mM L-Glutamine and 3.4 g/L D-(+)-Glucose. The pH of the medium is 7.0-7.4. Selection is 20 maintained by using 20 nM methotrexate (MTX), a competitive inhibitor of DHFR. 1 mL of an 20 µM MTX working solution is added to 1 L of complete medium.

No antibiotics are used. Samples for mycoplasma test are obtained periodically and analysed by the Mycoplasma Laboratory at the National Veterinary Institute, Uppsala, 25 Sweden. All samples tested were found to be negative for mycoplasma.

T-flasks

T-flasks (175 cm²) with 60 mL medium are seeded with 2.5-3 x 10⁵ cells/mL and incubated at 37 °C and 5 % CO₂ in a humidified incubator for 72-96 hours. After 3-4 days, a cell 30 concentration of 1-1.2 x10⁶ cells/mL is reached. The T-flasks are gently tapped to break any cell aggregates. The cell suspensions are centrifuged (110g, 8 min) in sterile tubes in a swing-out centrifuge (Eppendorf) at 20 °C. The collected supernatants are filtered (0.45 µm) and passed on to the purification process or stored frozen (-20 °C) for later use. Fresh and filtered supernatants from T-flasks hold a LAMAN activity of 30 mU/mL. The cell pellets 35 are resuspended in fresh medium at RT to a suitable cell density and passaged to new T-flasks cultures.

Counting of viable and dead cells is performed by using a haemocytometer and the trypan blue exclusion technique. Stored cells are frozen in liquid nitrogen in a working cell bank. New vials of cells are thawed regularly, at least every fifth month, and the LAMAN production is checked every week during the cultivation process.

5

Example 5: Partially developed purification scheme, formulation, filling and lyophilization

A purification process for rhLAMAN in 20-200 ml scale is developed for scale-up to large scale production. The quality and purity of the final product (rhLAMAN) is very high and according to the specifications (approved for clinical trials). The process includes a capture
10 step, 1-2 intermediate purification steps, 1 polishing step, 1-2 virus removal steps and 1 formulation step. 1 or more buffer exchange steps are also included (dia-filtration). The small-scale process is transferred to intermediate and finally large-scale production.

Experimental design: Several different chromatography gels are tested and performance of
15 the different steps will be analysed by a battery of analytical methods described briefly below:

<u>Enzyme activity:</u>	Alpha-mannosidase assay
<u>Total protein concentration:</u>	BCA analysis
20 <u>rhLAMAN concentration:</u>	rhLAMAN ELISA
<u>Purity:</u>	HPLC and SDS-PAGE
<u>Identity:</u>	HPLC
<u>HCP proteins:</u>	ELISA
<u>Endotoxin level:</u>	Outsource to contract Lab

25

Outline of purification process in 20 ml column scale

Step 1: Concentration/Diafiltration

Media produced in T-500 flasks (0.06U/ml) with expressed rhLAMAN is concentrated
30 approximately 10x for using Tangential flow filtration (TFF) against a Pellicon Biomax polysulphone filter with 100 kDa cut off.

Example: 0.5 – 1.0 litre spinner produced media is concentrated using TFF against a 50 cm² Biomax 100 kDa filter to 50 - 100 ml. Transmembrane pressure (TMP) is 25 (Pin = 30
35 psi; Pout = 20 psi).

WO 02/099092

PCT/DK02/00388

36

After that, diafiltration against 7 volumes of 20 mM Tris-HCl, pH 7.6 with TMP of 25 is applied. Specific activity of concentrated sample is 0.5 – 1.5 U/mg. Yield 70 – 90%

Step 2: Capture step – DEAE sepharose FF

- 5 Concentrated sample from step 1 is applied on a 20 ml DEAE sepharose packed in a 16 mm diameter column (Pharmacia XK 16) equilibrated with 20 mM Tris-HCl pH 7.6 (referred to as: standard buffer). Flow rate is 3 ml/min. Protein bound to the DEAE gel is then washed with 2-4 column volumes (CV) of standard buffer followed by 2-4 CV's of 30 mM NaCl in standard buffer.
- 10 rhLAMAN is eluted with a linear gradient from 30 – 300 mM NaCl in standard buffer for 20 minutes. Alternatively for the large scale production: Elution is achieved by applying 75 – 150 mM NaCl in standard buffer. The DEAE gel is washed with 2-4 CV's of 0.5 M NaCl in standard buffer followed by 2.4 CV's of 1.0 M NaOH (contact time 40 – 60 minutes).
- 15 Fractions containing rhLAMAN activity are pooled (specific activity: 2 – 5 U/mg) and used for further purification. Yield 70 – 90 %.

Step 3: Intermediate step 1

- 20 1. *Hydrophobic interaction chromatography: butyl, phenyl or octyl sepharose FF.*
Sample pool from step 2 is mixed 1:1 with 2.0 M Na₂SO₄ and applied on a 20 ml HIC column (butyl-, phenyl- or octyl-sepharose) packed in a 16 mm diameter column (Pharmacia XK 16) equilibrated with standard buffer + 1.0 M Na₂SO₄. Flow rate is 3-4 ml/min. Column is washed with 2-4 CV's of the same buffer. rhLAMAN
- 25 is eluted with standard buffer and fractions containing activity are pooled and used for further purification.
2. *Macroprep, Ceramic hydroxyapatite type I or II.*
Brief description: Equilibrate column (Gel volume = 20 ml) with 4-6 CV's of 10 mM Sodium phosphate H 7.6 (Buffer A). Flow rate 2-5 ml/min. Buffer exchange sample
- 30 from step 3 with TFF to Buffer A. Load sample from step 3 on column. Wash with 2-4 CV's of Buffer A. Elute with buffer A containing 100 mM NaCl. Collect peak containing rhLAMAN activity.

35

WO 02/099092

PCT/DK02/00388

37

Step 4: Intermediate step 2

1. *Macro prep, Ceramic hydroxyapatite type I or II, 40 um.*

Brief description: see Step 3

5

2. *Source 15 Q – anioexchanger*

Equilibrate column with 4 CV's of 200 mM Tris-HCl pH 7.6. Change to 5 CV's of 20 mM Tris-HCl pH 7.6 (standard buffer). Flow rate: 2- 5 ml/min. Load sample from step 3 (should be in standard buffer before application) onto column. Wash with 2- 4 CV's of standard buffer. Apply a shallow gradient from 0 to 100% of 1 M NaCl in standard buffer (flow rate 2ml/min, gradient time 50 minutes). Collect fractions containing rhLAMAN activity.

10

3. *Source 15 S – cation exchanger*

15

Should run in acidic pH. Equilibration buffer = 20 mM Sodium Acetate pH 4.5. Elute with a NaCl (increasing salt concentration) or pH (increasing pH) gradient.

Step 5: Polishing step

20

1. *Macro prep, Ceramic hydroxyapatite type II, 40 um.*

Description of parameters: see Step 3.

2. *Source 15 Q – anion exchanger*

Description of parameters: see Step 4.

25

3. *Source 15 S – cation exchanger*

Description of parameters: see Step 4.

30

4. *Affinity chromatography*

Step 6: Diafiltration / Formulation step

Tangential flow filtration (TFF) against a Pellicon Biomax polysulphone filter with 100 kDa cut off against 5-10 x Volumes of formulation buffer is performed. Two formulation buffers are tested:

35

Formulation buffer 1.

Na₂HPO₄ 3.10-3.50 mM

WO 02/099092

PCT/DK02/00388

38

NaH₂PO₄ 0.4-0.6 mM
Glycine 25-30 mM
Mannitol 220-250 mM
Water for injection (WFI)

5

Formulation buffer 2.

Sodium Citrate 4.0-5.0 mM
Citric Acid 0.3-0.8 mM
Mannitol 200-250 mM

10 Tween 80 3.0-5.0 mM

Water for injection (WFI)

The pH and osmolality in both Formulation buffers are balanced to 7.5 ± 0.2 and 300 ± 50 mOsm/kg respectively. Final protein concentration is according to the specification (>5
15 mg/ml).

Step 7: Formulation, Filling and Freeze-drying

Formulation and dosage form

In the development of dosage form the stability of rhLAMAN is focused. The development
20 process starts with an aqueous solution and will, most likely, end up as a freeze-dried
product. Two different formulations are tested: Formulation buffer 1 and Formulation
buffer 6, see Step 6.

Both these formulations are known to stabilize proteins in aqueous solutions as well as in
25 freeze-dried powders. The pH and osmolality in both Formulation buffers will be balanced
to 7.5 ± 0.2 and 300 ± 50 mOsm/kg respectively. Final protein concentration should be
according to the specification and in the range 4-10 mg/ml.

A freeze-dried product of rhLAMAN will be produced at a production unit according to EU
30 GMP practice. The filling and freeze-drying will be performed in a room classified as Class
A. During production the filling zone is monitored with particle count and settle plates. The
personnel are regularly trained according to EU GMP and monitored after each production
with glove prints. The sterility of equipment and materials are secured by validated
sterilization procedures.

35

Filling

The bulk drug substance of rhLAMAN are aseptically filled in sterile type I glass vials.

WO 02/099092

PCT/DK02/00388

39

Freeze-drying

The vials are freeze-dried with freeze-drying cycles specifically developed for rhLAMAN in the two different formulations described above. Nitrogen gas is filled into the vials in the
5 end of the cycle and eventually closed with stoppers and capped. The batch is finally analyzed and released according to the specification.

REFERENCES

10

Aebischer P, Goddard M, Signore AP, Timpson RL. 1994. Functional recovery in hemiparkinsonian primates transplanted with polymer-encapsulated PC12 cells. *Exp Neurol* 126:151-158

15 Aebischer P, Schluep M, Deglon N, Joseph JM, Hirt L, Heyd B, Goddard M, Hammang JP, Zurn AD, Kato AC, Regli F, Baetge EE. 1996. Intrathecal delivery of CNTF using encapsulated genetically modified xenogeneic cells in amyotrophic lateral sclerosis patients. *Nat Med* 2:696-699

20 Austin J, McAfee D, Armstrong D, O'Rourke M, Shearer L, Bachhawat B. 1964. Abnormal sulphatase activities in two human diseases (metachromatic leucodystrophy and gargoylism). *Biochem J* 93:15C-17C

Barth ML, Fensom A, Harris A. 1995. Identification of seven novel mutations associated
25 with metachromatic leukodystrophy. *Hum Mutat* 6(2):170-176

Bayever E, Ladisch S, Philippart M, Brill N, Nuwer M, Sparkes RS, Feig SA. 1985. Bone-marrow transplantation for metachromatic leucodystrophy. *Lancet* 2(8453):471-473

30 Bradley RT, Manowitz P. 1988. Electrochemical determination of lysosomal alpha-mannosidase activity using high-performance liquid chromatography. *Anal Biochem* 173(1):33-38

Braulke T, Hille A, Huttner WB, Hasilik A, von Figura K. 1987. Sulfated oligosaccharides in
35 human lysosomal enzymes. *Biochem Biophys Res Commun* 143(1):178-185

WO 02/099092

PCT/DK02/00388

40

Braulke T, Tippmer S, Chao HJ, von Figura K. 1990. Insulin-like growth factors I and II stimulate endocytosis but do not affect sorting of lysosomal enzymes in human fibroblasts. *J Biol Chem* 265(12):6650-6655

- 5 Deglon N, Heyd B, Tan SA, Joseph JM, Zurn AD, Aebischer P. 1996. Central nervous system delivery of recombinant ciliary neurotrophic factor by polymer encapsulated differentiated C2C12 myoblasts. *Hum Gene Ther* 7(17):2135-2146

- Dierks T, Schmidt B, von Figura K. 1997. Conversion of cysteine to formylglycine: a protein
10 modification in the endoplasmic reticulum. *Proc Natl Acad Sci USA* 94(22):11963-11968

- Draghia R, Letourneur F, Drugan C, Manicom J, Blanchot C, Kahn A, Poenaru L, Caillaud C. 1997. Metachromatic leukodystrophy: identification of the first deletion in exon 1 and of
15 nine novel point mutations in the lysosomal alpha-mannosidase gene. *Hum Mutat* 9(3):234-242

- Draper RK, Fiskum GM, Edmond J. 1976. Purification, molecular weight, amino acid, and subunit composition of lysosomal alpha-mannosidase from human liver. *Arch Biochem*
20 *Biophys* 177(2):525-538

- Dubois G, Turpin JC, Baumann N. 1975. Arylsulfatases isoenzymes in metachromatic leucodystrophy/detection of a new variant by electrophoresis improvement of quantitative assay. *Biomedicine* 23(3):116-119
25

Eto Y, Tokoro T, Liebaers I, Vamos E. 1982. Biochemical characterization of neonatal multiple sulfatase deficient (MSD) disorder cultured skin fibroblasts. *Biochem Biophys Res Commun* 106(2):429-34

- 30 Farooqui AA. 1976. Purification and properties of human placenta arylsulphatase A. *Arch Int Physiol Biochim* 84(3):479-492

- Farrell DF, MacMartin MP, Clark AF. 1979. Multiple molecular forms of lysosomal alpha-mannosidase in different forms of metachromatic leukodystrophy (alpha-mannosidosis).
35 *Neurology* 29(1):16-20

WO 02/099092

PCT/DK02/00388

41

Fujii T, Kobayashi T, Honke K, Gasa S, Ishikawa M, Shimizu T, Makita A. 1992. Proteolytic processing of human lysosomal alpha-mannosidase. *Biochim Biophys Acta* 1122(1):93-98

- 5 Gieselmann V, Polten A, Kreysing J, Kappler J, Fluharty A, von Figura K. 1991. Molecular genetics of metachromatic leukodystrophy. *Dev Neurosci* 13:222-227

Gieselmann V, Zlotogora J, Harris A, Wenger DA, Morris CP. 1994. Molecular genetics of metachromatic leukodystrophy. *Hum Mutat* 4:233-42

10

Gieselmann V, Matzner U, Hess B, Lullmann-Rauch R, Coenen R, Hartmann D, D'Hooge R, DeDeyn P, Nagels G. 1998. Metachromatic leukodystrophy: molecular genetics and an animal model. *J Inherit Metab Dis* 21:564-574

- 15 Gniot-Szulzycka J. 1974. Some properties of highly purified arylsulphatase A from human placenta. *Acta Biochim Pol* 21(3):247-254

Gustavson KH, Hagberg B. 1971. The incidence and genetics of metachromatic leucodystrophy in northern Sweden. *Acta Paediatr Scand* 60(5):585-590

20

Hess B, Saftig P, Hartmann D, Coenen R, Lullmann-Rauch R, Goebel HH, Evers M, von Figura K, D'Hooge R, Nagels G, De Deyn P, Peters C, Gieselmann V. 1996. Phenotype of lysosomal alpha-mannosidase-deficient mice: relationship to human metachromatic leukodystrophy. *Proc Natl Acad Sci USA* 93(25):14821-14826

25

Hickey WF, Kimura H. 1988. Perivascular microglial cells of the CNS are bone marrow-derived and present antigen in vivo. *Science* 239(4837):290-2

Hickey WF, Hsu BL, Kimura H. 1991. T-lymphocyte entry into the central nervous system.

- 30 *J Neurosci Res* 28(2):254-60

Inoue H, Seyama Y, Yamashita S. 1986. Specific determination of lysosomal alpha-mannosidase activity. *Experientia* 15;42(1):33-35

- 35 James GT. 1979. Essential arginine residues in human liver lysosomal alpha-mannosidase. *Arch Biochem Biophys* 197(1):57-62

WO 02/099092

PCT/DK02/00388

42

James GT, Thach AB, Klassen L, Austin JH Studies in metachromatic leukodystrophy: XV. 1985. Purification of normal and mutant lysosomal alpha-mannosidase from human liver. *Life Sci* 37(25):2365-2371

- 5 Joss V, Rogers TR, Hugh-Jones K, Beilby B, Joshi R, Williamson S, Foroozanfar N, Riches P, Turner M, Benson PD, Hobbs JR. 1982 Exp Hematol 10, 52

Kelly BM, Yu CZ, Chang PL. 1989. Presence of a lysosomal enzyme, arylsulfatase-A, in the prelysosome-endosome compartments of human cultured fibroblasts. *Eur J Cell Biol*

10 48(1):71-78

Klein MA, Frigg R, Flechsig E, Raeber AJ, Kalinke U, Bluethmann H, Bootz F, Suter M, Zinkernagel RM, Aguzzi A. 1997. A crucial role for B cells in neuroinvasive scrapie. *Nature* 390(6661):687-690

15

Kreysing J, Polten A, Hess B, von Figura K, Menz K, Steiner F, Gieselmann V. 1994. Structure of the mouse lysosomal alpha-mannosidase gene and cDNA. *Genomics* 19(2):249-256

- 20 Kreysing J, von Figura K, Gieselmann V. 1990. Structure of the lysosomal alpha-mannosidase gene. *Eur J Biochem* 1990. Eur J Biochem. 191(3): 627-631

Krivit W, Shapiro E, Kennedy W, Lipton M, Lockman L, Smith S, Summers CG, Wenger DA, Tsai MY, Ramsay NK, et al. 1990. Treatment of late infantile metachromatic

- 25 leukodystrophy by bone marrow transplantation. *N Engl J Med* 322:28-32

Krivit W, Shapiro E, Hoogerbrugge PM, Moser HW. 1992. State of the art review. Bone marrow transplantation treatment for storage diseases. Keystone. January 23, 1992. *Bone Marrow Transplant* 10 Suppl 1:87-96

30

Laidler PM, Waheed A, Van Etten RL. 1985. Structural and immunochemical characterization of human urine lysosomal alpha-mannosidase purified by affinity chromatography. *Biochim Biophys Acta* 827(1):73-83

- 35 Lee GD, van Etten RL. 1975. Evidence of an essential histidine residue in rabbit liver aryl sulfatase A. *Arch Biochem Biophys* 171(2):424-434

WO 02/099092

PCT/DK02/00388

43

Luijten JAFM, Van Der Heijden MCM, Rijksen G, Staal GEJ. 1978. Purification and characterization of lysosomal alpha-mannosidase from human urine. *J Mol Med.* 1978, 3, 213

- 5 Manowitz P, Fine LV, Nora R, Chokroverty S, Nathan PE, Fazzaro JM. 1988. A new electrophoretic variant of lysosomal alpha-mannosidase. *Biochem Med Metab Biol* 39(1):117-120

- 10 Mould DL, Dawson AM, Rennle JC. 1970. Very early replication of scrapie in lymphocytic tissue. *Nature* 228(273):779-780

Ohashi T, Watabe K, Sato Y, Saito I, Barranger JA, Matalon R, Eto Y. 1996. Gene therapy for metachromatic leukodystrophy. *Acta Paediatr Jpn* 38(2):193-201

- 15 Oya Y, Nakayasu H, Fujita N, Suzuki K, Suzuki K. 1998. Pathological study of mice with total deficiency of sphingolipid activator proteins (SAP knockout mice). *Acta Neuropathol (Berl)* 96(1):29-40

- 20 Poretz RD, Yang RS, Canas B, Lackland H, Stein S, Manowitz P. 1992. The structural basis for the electrophoretic isoforms of normal and variant human platelet arylsulphatase A. *Biochem J* 287 (Pt 3):979-983

- Porter MT, Fluharty AL, Kihara H. 1971. Correction of abnormal cerebroside sulfate metabolism in cultured metachromatic leukodystrophy fibroblasts. *Science* 1971,172, 25 1263-1265

- 30 Sangalli A, Taveggia C, Salviati A, Wrabetz L, Bordinon C, Severini GM. 1998. Transduced fibroblasts and metachromatic leukodystrophy lymphocytes transfer lysosomal alpha-mannosidase to myelinating glia and deficient cells in vitro. *Hum Gene Ther* 9(14):2111-2119

- Sarafian TA, Tsay KK, Jackson WE, Fluharty AL, Kihara H. 1985. Studies on the charge isomers of lysosomal alpha-mannosidase. *Biochem Med* 33(3):372-380 35

Schmidt B, Selmer T, Ingendoh A, von Figura K. 1995. A novel amino acid modification in sulfatases that is defective in multiple sulfatase deficiency. *Cell* 82(2):271-278

WO 02/099092

PCT/DK02/00388

44

Selmer T, Hallmann A, Schmidt B, Sumper M, von Figura K. 1996. The evolutionary conservation of a novel protein modification, the conversion of cysteine to serinesemialdehyde in arylsulfatase from *Volvox carteri*. *Eur J Biochem* 238:341-345

5

Shapira E, Nadler HL. 1975. Purification and some properties of soluble human liver arylsulfatases. *Arch Biochem Biophys* 170(1):179-187

Shapiro EG, Lockman LA, Balthazor M, Krivit W. 1995. Neuropsychological outcomes of several storage diseases with and without bone marrow transplantation. *J Inherit Metab Dis* 18:413-429

Stein C, Gieselmann V, Kreysing J, Schmidt B, Pohlmann R, Waheed A, Meyer HE, O'Brien JS, von Figura K. 1989. Cloning and expression of human lysosomal alpha-mannosidase. *J Biol Chem* 264:1252-1259

Stevens RL, Fluharty AL, Skokut MH, Kihara H. 1975. Purification and properties of arylsulfatase. A from human urine. *J Biol Chem* 250(7):2495-2501

20 Stevens RL, Fluharty AL, Killgrove AR, Kihara H. 1976. Microheterogeneity of lysosomal alpha-mannosidase from human tissues. *Biochim Biophys Acta* 445(3):661-671

von Figura K, Steckel F, Conary J, Hasilik A, Shaw E. 1986. Heterogeneity in late-onset metachromatic leukodystrophy. Effect of inhibitors of cysteine proteinases. *Am J Hum Genet*, 39, 371-382

Waheed A, Hasilik A, von Figura K. 1982. Synthesis and processing of lysosomal alpha-mannosidase in human skin fibroblasts. *Hoppe Seylers Z Physiol Chem* 363(4):425-430

30 Waheed A, van Etten RL. 1985. Phosphorylation and sulfation of lysosomal alpha-mannosidase accompanies biosynthesis of the enzyme in normal and carcinoma cell lines. *Biochim Biophys Acta* 847(1):53-61

WO 02/099092

45

PCT/DK02/00388

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>9</u> , line <u>13-15</u> .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH	
Address of depositary institution (including postal code and country) Maschroderweg 1b D-38124 Braunschweig Federal Republic of Germany	
Date of deposit 6 June 2002	Accession Number DSM ACC2549
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
As regards the respective Patent Offices of the respective designated states, the applicant requests that a sample of the deposited microorganism only be made available to an expert nominated by the requester until the date on which the patent is granted or the date on which the application has been refused or withdrawn or is deemed to be withdrawn.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau (specify the general nature of the indications e.g. "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer
SUBSTITUTE SHEET	

WO 02/099092

PCT/DK02/00388

46

CLAIMS

1. A cell capable of producing recombinant human LAMAN (rhLAMAN), said cell comprising the 3066 basepair EcoRI - XbaI fragment of a human cDNA which codes for a human
5 LAMAN protein in which the position corresponding to position 186 of the full length hLAMAN protein is Aspartic acid (Asp, D).
2. A cell according to claim 1 comprising a DNA sequence which codes for the amino acid sequence shown in SEQ ID NO 12.
- 10 3. A cell according to claim 1 obtained by insertion of a DNA fragment comprising the 3066 basepair EcoRI - XbaI fragment of the DNA fragment having the sequence shown in SEQ ID NO 2.
- 15 4. A cell according to any of claims 1 - 3 obtained by use of the expression plasmid pLamanExp1 having SEQ ID NO. 2.
5. A cell according to any of claims 1 - 4 obtained by transfection of a non-human mammalian cell line.
- 20 6. A cell according to claim 5 obtained by transfection of chinese hamster ovary (CHO) cells.
7. A cell according to any of claims 1 - 6 obtained by the culture of the human LAMAN
25 production cell line CHO DG44.42 subclone 20L-31.62.5 which has been deposited at the DSMZ for the purposes of patent deposit according to the Budapest Treaty on 6 June 2002.
8. A method for the preparation of recombinant human LAMAN, the method comprising
30 a) introducing, into a suitable vector, a nucleic acid fragment comprising a DNA fragment which codes for the amino acid sequence shown SEQ ID NO 12;
b) transforming a cell with the vector obtained in step a);
35 c) culturing the transformed host cell under conditions facilitating expression of the nucleic acid sequence;
d) recovering the expression product from the culture.

WO 02/099092

PCT/DK02/00388

47

9. A method according to claim 8 further comprising a fermentation step.

10. A method according to claim 8 or 9 further comprising a purification step.

5 11. A method according to any of claims 8-10, wherein the nucleic acid fragment comprises the 3066 basepair EcoRI - XbaI fragment of the DNA fragment shown in SEQ ID NO 2.

12. An expression plasmid pLamanExp1 having the sequence shown in SEQ ID NO. 2.

10

13. A rhLAMAN produced by the method of any of claims 8-11.

14. Use of the rhLAMAN according to claim 13 for the preparation of a medicament for the treatment of alfa mannosidosis.

15

1/9




Fig. 1

2/9

STEP I




Fig. 2

3/9

STEP II




Fig. 2 continued

4/9




Fig. 3

5/9




Fig. 4

WO 02/099092

PCT/DK02/00388

6/9




Fig. 5

WO 02/099092

PCT/DK02/00388

7/9




Fig. 6

WO 02/099092

PCT/DK02/00388

8/9




Fig. 7

WO 02/099092

PCT/DK02/00388

9/9




Fig. 8

WO 02/099092

1/13

PCT/DK02/00388

SEQUENCE LISTING

<110> Jens Fogh
Meher Irani
Claes Andersson
Cecilia Weigelt

<120> Production of recombinant human
lysosomal alpha-mannosidase.

<130> P25378PC01

<150> US 60/296.142

<151> 2001-06-07

<160> 12

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 6591

<212> DNA

<213> Artificial Sequence

<220>

<223> Expression plasmid pAsaExp1

<400> 1

```

agatcttcaa tattggccat tagccatatt attcattggt tatatagcat aaatcaatat 60
tggctattgg ccattgcata cgttgtatct atatcataat atgtacattt atattggctc 120
atgtccaata tgaccgccat gttggcattg attattgact agttattaat agtaatcaat 180
tacggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa 240
tggccgcgct ggctgaccgc ccaacgaccc cgcgccattg acgtcaataa tgacgtatgt 300
tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacgta 360
aactgcccac ttggcagtac atcaagtgtat tcatatgccca agtccgcccc ctattgacgt 420
caatgacggt aaatggcccc cctggcatta tgcccagtac atgaccttac gggactttcc 480
tacttggcag tacatctacg tattagtcat cgctattacc atggtgatgc ggttttgga 540
gtacaccaat gggcggtgat agcggtttga ctacgaggga tttccaagtc tccaccccat 600
tgacgtcaat gggagtttgt tttggcacca aaatcaacgg gactttccaa aatgtcgtaa 660
caactgcgat cgcgcgcccc gttgacgcaa atgggcggta ggcgtgtacg gtgggaggtc 720
tatataagca gagctcggtt agtgaaccgt cagatcacta gaagctttat tgcggtagt 780
tatcacagtt aaattgctaa cgcagtcagt gcttctgaca caacagtcctc gaacttaagc 840
tgacgtgact ctcttaaggt agccttgtag aagttggtcg tgaggcactg ggcaggtaag 900
tatcaaggtt acaagacagg ttaaggaga ccaatagaaa ctgggcttgt cgagacagag 960
aagactcttg cgtttctgat aggcacctat tggctctact gacatccact ttgcctttct 1020
ctccacaggt gtccactccc agttcaatta cagctcttaa ggctagagta cttaatacga 1080
ctcactatag gctagcctcg agaattcgcc accatggggg caccgcggtc cctcctcctg 1140
gccttggtcg ctggcctggc cgttgacagt ccgcccaaca tcgtgctgat ctttgcggac 1200
gacctcggtc atggggacct gggctgctat gggcacccca gctctaccac tcccaacctg 1260
gaccagcttg cggcgggagg gctgcggttc acagacttct acgtgcctgt gtctctgtgc 1320
acacctcta gggccgcccc cctgaccggc cggtcccggt ttcggatggg catgtaccct 1380
ggcgctcctg tgcccagctc ccgggggggc ctgcccctgg aggaggtgac cgtggccgaa 1440
gtcctggctg cccgaggcta cctcacagga atggccggca agtggcacct tggggtgggg 1500
cctgaggggg ccttctgccc ccccatcag ggcttccatc gatttctagg catcccgta 1560
tccacagacc agggccccct ccagaacctg acctgcttcc cgccggccac tccttgcgac 1620
ggtggctgtg accagggcct ggtccccatc ccaactgttg ccaacctgtc cgtggaggcg 1680
cagccccctt ggctgcccgg actagaggcc cgctacatgg ctttcgccc tgacctcatg 1740
ccgacgcccc agcgccagga tcgccccttc ttctgtact atgcctctca ccacaccac 1800

```

WO 02/099092

2/13

PCT/DK02/00388

taccctcagt	tcagtgggca	gagctttgca	gagcgttcag	gcgcggggcc	atttggggac	1860
tccctgatgg	agctggatgc	agctgtgggg	accctgatga	cagccatagg	ggacctgggg	1920
ctgcttgaag	agacgctggt	catcttctact	gcagacaatg	gacctgagac	catgctgatg	1980
tcccgaggcg	gctgctccgg	tctcttgccg	tgtggaaagg	gaacgacct	cgaggggcgt	2040
gtccgagagc	ctgccttgcc	cttctggcca	ggcatatcg	ctcccgccgt	gaccacagag	2100
ctggccagct	ccctggacct	gctgcctacc	ctggcagccc	tggctggggc	cccactgccc	2160
aatgtcacct	tggatggctt	tgacctcagc	cccctgctgc	tgggcacagg	caagagccct	2220
cggcagtcct	tcttcttcta	cccgctcctac	ccagacgagg	tccgtggggg	ttttgctgtg	2280
cggactggaa	agtacaaggc	tcacttcttc	accaggggt	ctgcccacag	tgataccact	2340
gcagaccctg	cctgccacgc	ctccagctct	ctgactgctc	atgagccccc	gctgctctat	2400
gacctgtcca	aggaccctgg	tgagaactac	aacctgctgg	gggggtgtggc	cgggggccacc	2460
ccagaggtgc	tgaagccct	gaaacagctt	cagctgctca	aggcccagtt	agacgcagct	2520
gtgaccttcg	gccccagcca	ggtggcccg	ggcgaggacc	ccgccctgca	gatctgctgt	2580
catctggct	gcaccccccg	cccagcttgc	tgccattgcc	cagatcccca	tgccctgagg	2640
cccctcggt	ggcctgggca	tgtgatggct	cctcactggt	ctagagtcca	cccggggcgc	2700
cgcttccctt	tagtgagggt	taatgcttcg	agcagacatg	ataagataca	ttgatgagtt	2760
tggaacaacc	acaactagaa	tgcaagtcaa	aaaatgcttt	atttgtgaaa	tttgtgatgt	2820
tattgcttta	tttgtaacca	ttataagctg	caataaaca	gttaacaaca	acaattgcat	2880
tcattttatg	tttcagggtt	agggggagat	gtgggaggtt	ttttaagca	agtaaacct	2940
ctacaaatgt	ggtaaaatcc	gataaggatc	gatccgggct	ggcgtaatag	cgaagaggcc	3000
cgaccgcatc	gcccttccca	acagttgcgc	agcctgaatg	gcgaatggac	gcgccctgta	3060
gcggcgcat	aagcgcgcg	ggtgtggtg	ttacgcgcag	cgtgaccgct	acacttgcca	3120
gcgcctagc	gcccgtcct	ttcgctttct	tcccttctt	tctcgccacg	ttcgccggct	3180
ttcccgtca	agctctaaat	cgggggctcc	ctttagggtt	ccgatttaga	gctttacgat	3240
acctcgaccg	caaaaaactt	gatttgggtg	atggttcacg	tagtgggcca	tcgccctgat	3300
agacggtttt	tcgccctttg	acgttggagt	ccacgttctt	taatagtga	ctcttgttcc	3360
aaactggaac	aacactcaac	cctatctcgg	tctattcttt	tgatttataa	gggattttgg	3420
ggatttccgc	ctattggtta	aaaaatgagc	tgatttaaca	aaaatttaac	gcgaattaat	3480
tctgtggaat	gtgtgtcagt	tagggtgtgg	aaagtcccca	ggctccccag	gcaggcagaa	3540
gtatgcaaag	catgcatctc	aattagtcag	caaccagggt	tgaaaagtcc	ccaggctccc	3600
cagcaggcag	aagtatgcaa	agcatgcac	tcaattagtc	agcaaccata	gtcccgcgcc	3660
taactccgcc	catcccgccc	ctaactccgc	ccagttccgc	ccattctccg	ccccatggct	3720
gactaatttt	ttttatttat	gcagaggccg	aggccgcctc	tgccctctag	ctattccaga	3780
agtagtgagg	aggctttttt	ggaggccctg	gcttttgcaa	aaagctccc	ggatggttcg	3840
accattgaac	tgcatcgtcg	ccgtgtccca	aaatatgggg	attggcaaga	acggagacct	3900
accctggcct	ccgctcagga	acgagttcaa	gtacttccaa	agaatgacca	caacctcttc	3960
agtggaaggt	aaacagaatc	tggtgattat	gggtaggaaa	acctgggtct	ccattcctga	4020
gaagaatcga	cccttaagg	acagaattaa	ttatagttct	agtagagaac	tcaagaaac	4080
accacgagga	gctcattttc	ttgccaaaag	tttgatgat	gccttaagac	ttattgaaca	4140
accggaattg	gcaagtaaa	tagacatggt	ttggatagtc	ggaggcagtt	ctgtttacca	4200
ggaagccatg	aatcaaccag	gccaccttag	actctttgtg	acaaggatca	tgagggaatt	4260
tgaaagtgc	acgtttttcc	cagaaaattga	tttggggaaa	tataaacttc	tcccagaata	4320
cccaggcgct	ctctctgagg	tccaggagga	aaaaggcatc	aagtataagt	ttgaagtcta	4380
cgagaagaaa	gactaattcg	aaatgaccga	ccaagcgacg	cccaacctgc	catcacgatg	4440
gcgcgaataa	aatatcttta	ttttcattac	atctgtgtgt	tggttttttg	tgtgaatcga	4500
tagcgataag	gatccgcgta	tgtgtcactc	tcagtacaat	ctgctctgat	gccgcatagt	4560
taagccagcc	ccgacacccg	ccaacacccg	ctgacgcgcc	ctgacgggct	tgtctgctcc	4620
cggcatccgc	ttacagacaa	gctgtgaccg	tctccgggag	ctgcatgtgt	cagaggtttt	4680
caccgtcatc	accgaaacgc	gcgagacgaa	agggcctcgt	gatacgccct	tttttatagg	4740
ttaatgtcat	gataataatg	gtttcttaga	cgtcagggtg	cacttttctg	ggaaatgtgc	4800
gcggaacccc	tatttgttta	tttttctaaa	tacattcaaa	tatgtatccg	ctcatgagac	4860
aataaccctg	ataaatgctt	caataatatt	gaaaaaggaa	gagtatgagt	attcaacatt	4920
tccgtgtcgc	ccttattccc	ttttttcgcc	cattttgcct	tcctgttttt	gctcaccag	4980
aaacgctggt	gaaagtaaaa	gatgctgaag	atcagttggg	tgacagagt	ggttacatcg	5040
aactggatct	caacagcggt	aagatccttg	agagttttcg	ccccgaagaa	cgttttccaa	5100
tgatgagcac	ttttaaagtt	ctgctatgtg	gcgcggtatt	atcccgtatt	gacgcggg	5160
aagagcaact	cggtgcgcgc	atacactatt	ctcagaatga	cttggttgag	tactcaccag	5220
tcacagaaaa	gcattcttacg	gatggcatga	cagtaagaga	attatgcagt	gctgccataa	5280
ccatgagtga	taacactgcg	gccaacttac	ttctgacaac	gatcgaggga	ccgaaggagc	5340

WO 02/099092

3/13

PCT/DK02/00388

taaccgcttt tttgcacaac atggggggtc atgtaactcg ccttgatcgt tgggaaccgg 5400
agctgaatga agccatacca aacgacgagc gtgacaccac gatgcctgta gcaatggcaa 5460
caacgttgcg caaactatta actggcgaaac tacttactct agcttcccgg caacaattaa 5520
tagactggat ggaggcggt aaagtgcag gaccacttct gcgctcgcc cttccggctg 5580
gctggtttat tgctgataaa tctggagccg gtgagcgtgg gtctcgcggt atcattgcag 5640
cactggggcc agatggtaag ccctcccgtc tcgtagtatt ctacacgacg gggagtcagg 5700
caactatgga tgaacgaaat agacagatcg ctgagatagg tgcctcactg attaagcatt 5760
ggtaactgtc agaccaagtt tactcatata tacttttagt tgatttaaaa cttcattttt 5820
aattttaaag gatctagggt aagatccttt ttgataatct catgaccaa atcccttaac 5880
gtgagttttc gttccactga gcgtcagacc ccgtagaaaa gatcaaagga tcttcttgag 5940
atcctttttt tctgcgcgta atctgctgct tgcaaacaaa aaaaccaccg ctaccagcgg 6000
tggtttgttt gccggatcaa gagctaccaa ctctttttcc gaaggtaact ggcttcagca 6060
gagcgcagat accaaatact gtccttctag tgtagccgta gttaggccac cacttcaaga 6120
actctgtagc accgcctaca tacctcgctc tgctaactct gttaccagtg gctgctgcca 6180
gtggcgataa gtcgtgtctt accgggttgg actcaagacg atagttaccg gataaggcgc 6240
agcggtcggg ctgaacgggg ggttcgtgca cacagcccag cttggagcga acgacctaca 6300
ccgaactgag atacctacag cgtgagctat gagaaagcgc cacgcttccc gaaggagaaa 6360
aggcggacag gtatccggta agcggcaggg tcggaacagg agagcgacg agggagcttc 6420
cagggggaaa cgcctggtat ctttatagtc ctgtcgggtt tcgccacctc tgacttgagc 6480
gtcgattttt gtgatgctcg tcaggggggc ggagcctatg gaaaaacgcc agcaacgcgg 6540
cctttttacg gttcctggcc ttttgctggc cttttgctca catggctcga c 6591

<210> 2
<211> 8079
<212> DNA
<213> Artificial Sequence

<220>
<223> Expression plasmid pLamanExp1

<221> CDS
<222> (1114)...(4149)
<223> LAMAN coding region

<221> misc feature
<222> (1669)...(1671)
<223> The Asp codon, corresponding to position 186 in the amino acid sequence.

<400> 2
agatcttcaa tattggccat tagccatatt attcattggt tatatagcat aaatcaatat 60
tggtatttgg ccattgcata cgttgtatct atatcataat atgtacattt atattggctc 120
atgtccaata tgaccgccat gttggcattg attattgact agttattaat agtaatcaat 180
tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa 240
tgcccgcctt ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt 300
tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggtg 360
aactgcccac ttggcagtag atcaagtgtc tcatatgcca agtccgcccc ctattgacgt 420
caatgacggt aaatggcccc cctggcatta tgcccagtag atgaccttac gggactttcc 480
tacttggcag tacatctacg tatttagtcat cgctattacc atgggtgatg gggttttgca 540
gtacaccaat gggcgtggat agcgggttga ctacgggga tttccaagtc tccaccccat 600
tgacgtcaat ggggctttgt tttggcacca aaatcaacgg gactttccaa aatgtcgtaa 660
caactgcgat cgcgcgcccc gttgacgcaa atgggcggta ggcgtgtacg gtgggaggtc 720
tatataagca gagctcgttt agtgaaccgt cagatcacta gaagctttat tgcggtagtt 780
tatcacagtt aaattgctaa cgcagtcagt gcttctgaca caacagtctc gaacttaagc 840
tgacgtgact ctcttaaggt agccttgtag aagttggtcg tgaggcactg ggcaggttaag 900
tatcaagggtt acaagacagg tttaaggaga ccaatagaaa ctgggcttgt cgagacagag 960
aagactcttg cgtttctgat aggcacctat tggcttact gacatccact ttgcctttct 1020
ctccacaggt gtccactccc agttcaatta cagctcttaa ggctagagta cttaatagca 1080
ctcactatag gctagcctcg agaattcgcc gcc atg ggc gcc tac gcg cgg gct 1134

PCT/DK02/00388

Met Gly Ala Tyr Ala Arg Ala
1 5

ttc ttc ttt ggg cgc ctt gat tat caa gat aag tgg gta cgg atg cag 1806
Phe Phe Phe Gly Arg Leu Asp Tyr Gln Asp Lys Trp Val Arg Met Gln
220 225 230

WO 02/099092

PCT/DK02/00388

5/13

aag ctg gag atg gag cag gtg tgg cgg gcc agc acc agc ctg aag ccc	1854
Lys Leu Glu Met Glu Gln Val Trp Arg Ala Ser Thr Ser Leu Lys Pro	
235 240 245	
ccg acc gcg gac ctc ttc act ggt gtg ctt ccc aat ggt tac aac ccg	1902
Pro Thr Ala Asp Leu Phe Thr Gly Val Leu Pro Asn Gly Tyr Asn Pro	
250 255 260	
cca agg aat ctg tgc tgg gat gtg ctg tgt gtc gat cag ccg ctg gtg	1950
Pro Arg Asn Leu Cys Trp Asp Val Leu Cys Val Asp Gln Pro Leu Val	
265 270 275	
gag gac cct cgc agc ccc gag tac aac gcc aag gag ctg gtc gat tac	1998
Glu Asp Pro Arg Ser Pro Glu Tyr Asn Ala Lys Glu Leu Val Asp Tyr	
280 285 290 295	
ttc cta aat gtg gcc act gcc cag ggc cgg tat tac cgc acc aac cac	2046
Phe Leu Asn Val Ala Thr Ala Gln Gly Arg Tyr Tyr Arg Thr Asn His	
300 305 310	
act gtg atg acc atg ggc tgc gac ttc caa tat gag aat gcc aac atg	2094
Thr Val Met Thr Met Gly Ser Asp Phe Gln Tyr Glu Asn Ala Asn Met	
315 320 325	
tgg ttc aag aac ctt gac aag ctc atc cgg ctg gta aat gcg cag cag	2142
Trp Phe Lys Asn Leu Asp Lys Leu Ile Arg Leu Val Asn Ala Gln Gln	
330 335 340	
gca aaa gga agc agt gtc cat gtt ctc tac tcc acc ccc gct tgt tac	2190
Ala Lys Gly Ser Ser Val His Val Leu Tyr Ser Thr Pro Ala Cys Tyr	
345 350 355	
ctc tgg gag ctg aac aag gcc aac ctc acc tgg tca gtg aaa cat gac	2238
Leu Trp Glu Leu Asn Lys Ala Asn Leu Thr Trp Ser Val Lys His Asp	
360 365 370 375	
gac ttc ttc cct tac gcg gat ggc ccc cac cag ttc tgg acc ggt tac	2286
Asp Phe Phe Pro Tyr Ala Asp Gly Pro His Gln Phe Trp Thr Gly Tyr	
380 385 390	
ttt tcc agt cgg ccg gcc ctc aaa cgc tac gag cgc ctc agc tac aac	2334
Phe Ser Ser Arg Pro Ala Leu Lys Arg Tyr Glu Arg Leu Ser Tyr Asn	
395 400 405	
ttc ctg cag gtg tgc aac cag ctg gag gcg ctg gtg ggc ctg gcg gcc	2382
Phe Leu Gln Val Cys Asn Gln Leu Glu Ala Leu Val Gly Leu Ala Ala	
410 415 420	
aac gtg gga ccc tat ggc tcc gga gac agt gca ccc ctc aat gag gcg	2430
Asn Val Gly Pro Tyr Gly Ser Gly Asp Ser Ala Pro Leu Asn Glu Ala	
425 430 435	
atg gct gtg ctc cag cat cac gac gcc gtc agc ggc acc tcc cgc cag	2478
Met Ala Val Leu Gln His His Asp Ala Val Ser Gly Thr Ser Arg Gln	
440 445 450 455	
cac gtg gcc aac gac tac gcg cgc cag ctt gcg gca ggc tgg ggg cct	2526
His Val Ala Asn Asp Tyr Ala Arg Gln Leu Ala Ala Gly Trp Gly Pro	
460 465 470	

WO 02/099092

6/13

PCT/DK02/00388

tgc gag gtt ctt ctg agc aac gcg ctg gcg cgg ctc aga ggc ttc aaa	2574
Cys Glu Val Leu Leu Ser Asn Ala Leu Ala Arg Leu Arg Gly Phe Lys	
475 480 485	
gat cac ttc acc ttt tgc caa cag cta aac atc agc atc tgc ccg ctc	2622
Asp His Phe Thr Phe Cys Gln Gln Leu Asn Ile Ser Ile Cys Pro Leu	
490 495 500	
agc cag acg gcg gcg cgc ttc cag gtc atc gtt tat aat ccc ctg ggg	2670
Ser Gln Thr Ala Ala Arg Phe Gln Val Ile Val Tyr Asn Pro Leu Gly	
505 510 515	
cgg aag gtg aat tgg atg gta cgg ctg ccg gtc agc gaa ggc gtt ttc	2718
Arg Lys Val Asn Trp Met Val Arg Leu Pro Val Ser Glu Gly Val Phe	
520 525 530 535	
gtt gtg aag gac ccc aat ggc agg aca gtg ccc agc gat gtg gta ata	2766
Val Val Lys Asp Pro Asn Gly Arg Thr Val Pro Ser Asp Val Val Ile	
540 545 550	
ttt ccc agc tca gac agc cag gcg cac cct ccg gag ctg ctg ttc tca	2814
Phe Pro Ser Ser Asp Ser Gln Ala His Pro Pro Glu Leu Leu Phe Ser	
555 560 565	
gcc tca ctg ccc gcc ctg ggc ttc agc acc tat tca gta gcc cag gtg	2862
Ala Ser Leu Pro Ala Leu Gly Phe Ser Thr Tyr Ser Val Ala Gln Val	
570 575 580	
cct cgc tgg aag ccc cag gcc cgc gca cca cag ccc atc ccc aga aga	2910
Pro Arg Trp Lys Pro Gln Ala Arg Ala Pro Gln Pro Ile Pro Arg Arg	
585 590 595	
tcc tgg tcc cct gct tta acc atc gaa aat gag cac atc cgg gca acg	2958
Ser Trp Ser Pro Ala Leu Thr Ile Glu Asn Glu His Ile Arg Ala Thr	
600 605 610 615	
ttt gat cct gac aca ggg ctg ttg atg gag att atg aac atg aat cag	3006
Phe Asp Pro Asp Thr Gly Leu Leu Met Glu Ile Met Asn Met Asn Gln	
620 625 630	
caa ctc ctg ctg cct gtt cgc cag acc ttc ttc tgg tac aac gcc agt	3054
Gln Leu Leu Leu Pro Val Arg Gln Thr Phe Phe Trp Tyr Asn Ala Ser	
635 640 645	
ata ggt gac aac gaa agt gac cag gcc tca ggt gcc tac atc ttc aga	3102
Ile Gly Asp Asn Glu Ser Asp Gln Ala Ser Gly Ala Tyr Ile Phe Arg	
650 655 660	
ccc aac caa cag aaa ccg ctg cct gtg agc cgc tgg gct cag atc cac	3150
Pro Asn Gln Gln Lys Pro Leu Pro Val Ser Arg Trp Ala Gln Ile His	
665 670 675	
ctg gtg aag aca ccc ttg gtg cag gag gtg cac cag aac ttc tca gct	3198
Leu Val Lys Thr Pro Leu Val Gln Glu Val His Gln Asn Phe Ser Ala	
680 685 690 695	
tgg tgt tcc cag gtg gtt cgc ctg tac cca gga cag cgg cac ctg gag	3246
Trp Cys Ser Gln Val Val Arg Leu Tyr Pro Gly Gln Arg His Leu Glu	

WO 02/099092

PCT/DK02/00388

7/13

700										705										710										
cta	gag	tgg	tcg	gtg	ggg	ccg	ata	cct	gtg	ggc	gac	acc	tgg	ggg	aag		3294													
Leu	Glu	Trp	Ser	Val	Gly	Pro	Ile	Pro	Val	Gly	Asp	Thr	Trp	Gly	Lys															
			715						720					725																
gag	gtc	atc	agc	cgt	ttt	gac	aca	ccg	ctg	gag	aca	aag	gga	cgc	ttc		3342													
Glu	Val	Ile	Ser	Arg	Phe	Asp	Thr	Pro	Leu	Glu	Thr	Lys	Gly	Arg	Phe															
			730				735						740																	
tac	aca	gac	agc	aat	ggc	cgg	gag	atc	ctg	gag	agg	agg	cgg	gat	tat		3390													
Tyr	Thr	Asp	Ser	Asn	Gly	Arg	Glu	Ile	Leu	Glu	Arg	Arg	Arg	Asp	Tyr															
			745				750						755																	
cga	ccc	acc	tgg	aaa	ctg	aac	cag	acg	gag	ccc	gtg	gca	gga	aac	tac		3438													
Arg	Pro	Thr	Trp	Lys	Leu	Asn	Gln	Thr	Glu	Pro	Val	Ala	Gly	Asn	Tyr															
			760				765				770				775															
tat	cca	gtc	aac	acc	cgg	att	tac	atc	acg	gat	gga	aac	atg	cag	ctg		3486													
Tyr	Pro	Val	Asn	Thr	Arg	Ile	Tyr	Ile	Thr	Asp	Gly	Asn	Met	Gln	Leu															
					780					785					790															
act	gtg	ctg	act	gac	cgc	tcc	cag	ggg	ggc	agc	agc	ctg	aga	gat	ggc		3534													
Thr	Val	Leu	Thr	Asp	Arg	Ser	Gln	Gly	Gly	Ser	Ser	Leu	Arg	Asp	Gly															
					795				800						805															
tcg	ctg	gag	ctc	atg	gtg	cac	cga	agg	ctg	ctg	aag	gac	gat	gga	cgc		3582													
Ser	Leu	Glu	Leu	Met	Val	His	Arg	Arg	Leu	Leu	Lys	Asp	Asp	Gly	Arg															
					810				815						820															
gga	gta	tcg	gag	cca	cta	atg	gag	aac	ggg	tcg	ggg	gcg	tgg	gtg	cga		3630													
Gly	Val	Ser	Glu	Pro	Leu	Met	Glu	Asn	Gly	Ser	Gly	Ala	Trp	Val	Arg															
					825				830						835															
ggg	cgc	cac	ctg	gtg	ctg	ctg	gac	aca	gcc	cag	gct	gca	gcc	gcc	gga		3678													
Gly	Arg	His	Leu	Val	Leu	Leu	Asp	Thr	Ala	Gln	Ala	Ala	Ala	Ala	Gly															
					840					845					850															
cac	cgg	ctc	ctg	gcg	gag	cag	gag	gtc	ctg	gcc	cct	cag	gtg	gtg	ctg		3726													
His	Arg	Leu	Leu	Ala	Glu	Gln	Glu	Val	Leu	Ala	Pro	Gln	Val	Val	Leu															
					860					865					870															
gcc	ccg	ggt	ggc	ggc	gcc	gcc	tac	aat	ctc	ggg	gct	cct	ccg	cgc	acg		3774													
Ala	Pro	Gly	Gly	Gly	Ala	Ala	Tyr	Asn	Leu	Gly	Ala	Pro	Pro	Arg	Thr															
					875				880						885															
cag	ttc	tca	ggg	ctg	cgc	agg	gac	ctg	ccg	ccc	tcg	gtg	cac	ctg	ctc		3822													
Gln	Phe	Ser	Gly	Leu	Arg	Arg	Asp	Leu	Pro	Pro	Ser	Val	His	Leu	Leu															
					890				895						900															
acg	ctg	gcc	agc	tgg	ggc	ccc	gaa	atg	gtg	ctg	ctg	cgc	ttg	gag	cac		3870													
Thr	Leu	Ala	Ser	Trp	Gly	Pro	Glu	Met	Val	Leu	Leu	Arg	Leu	Glu	His															
					905				910						915															
cag	ttt	gcc	gta	gga	gag	gat	tcc	gga	cgt	aac	ctg	agc	gcc	ccc	gtt		3918													
Gln	Phe	Ala	Val	Gly	Glu	Asp	Ser	Gly	Arg	Asn	Leu	Ser	Ala	Pro	Val															
					920					925					930															
acc	ttg	aac	ttg	agg	gac	ctg	ttc	tcc	acc	ttc	acc	atc	acc	cgc	ctg		3966													

PCT/DK02/00388

8/13

Thr	Leu	Asn	Leu	Arg	Asp	Leu	Phe	Ser	Thr	Phe	Thr	Ile	Thr	Arg	Leu		
				940					945				950				
cag	gag	acc	acg	ctg	gtg	gcc	aac	cag	ctc	cgc	gag	gca	gcc	tcc	agg	4014	
Gln	Glu	Thr	Thr	Leu	Val	Ala	Asn	Gln	Leu	Arg	Glu	Ala	Ala	Ser	Arg		
				955					960				965				
ctc	aag	tgg	aca	aca	aac	aca	ggc	ccc	aca	ccc	cac	caa	act	ccg	tac	4062	
Leu	Lys	Trp	Thr	Thr	Asn	Thr	Gly	Pro	Thr	Pro	His	Gln	Thr	Pro	Tyr		
				970					975				980 <td></td>				
cag	ctg	gac	ccg	gcc	aac	atc	acg	ctg	gaa	ccc	atg	gaa	atc	cgc	act	4110	
Gln	Leu	Asp	Pro	Ala	Asn	Ile	Thr	Leu	Glu	Pro	Met	Glu	Ile	Arg	Thr		
				985					990				995 <td></td>				
ttc	ctg	gcc	tca	gtt	caa	tgg	aag	gag	gtg	gat	ggg	tag	gtctgtctggg			4159	
Phe	Leu	Ala	Ser	Val	Gln	Trp	Lys	Glu	Val	Asp	Gly	*					
				1000					1005				1010				
atggggccctc	tagagtcgac	ccggggcgggc	gcttcccttt	agtgaggggtt	aatgcttcga	4219											
gcagacatga	taagatacat	tgatgagttt	ggacaaacca	caactagaat	gcagtgaaaa	4279											
aaatgcttta	tttgtgaaat	ttgtgatgct	attgctttat	ttgtaaccat	tataagctgc	4339											
aataaacaag	ttaacaacaa	caattgcatt	cattttatgt	ttcaggttca	gggggagatg	4399											
tgggaggttt	tttaaagcaa	gtaaaacctc	tacaaatgtg	gtaaaatccg	ataaggatcg	4459											
atccgggctg	gcgtaatagc	gaagaggccc	gcaccgatcg	cccttcccaa	cagttgcgca	4519											
gcctgaatgg	cgaatggacg	cgccctgtag	cgcgcgatta	agcgcggcgg	gtgtgggtgt	4579											
tacgcgcagc	gtgaccgcta	cacttggcag	cgccctagcg	cccgtctcct	tcgctttctt	4639											
cccttccctt	ctcgccacgt	tcgccggctt	tccccgtcaa	gctctaaatc	gggggctccc	4699											
ctttagggttc	cgatttagag	ctttacggca	cctcgaccgc	aaaaaacctg	atttgggtga	4759											
tgggttcacgt	agtgggccat	cgccctgata	gacggttttt	cgccctttga	cgttgagatc	4819											
cacgttcttt	aatagtggac	tcttgttcca	aactggaaca	acactcaacc	ctatctcggt	4879											
ctattctttt	gatttataag	ggattttggg	gatttcggcc	tattggttaa	aaaatgagct	4939											
gatttaacaa	aaatttaacg	cgaattaatt	ctgtggaatg	tgtgtcagtt	agggtgtgga	4999											
aagtccccag	gctccccagg	caggcagaag	tatgcaaagc	atgcatctca	attagtccgc	5059											
aaccagggtgt	ggaaagtccc	caggctcccc	agcaggcaga	agtatgcaa	gcatgcatct	5119											
caattagtca	gcaaccatag	tcccgcacct	caactcggcc	atcccccccc	taactccgcc	5179											
cagttccgcc	cattctccgc	cccatggctg	actaattttt	tttattttatg	cagaggccga	5239											
ggccgcctct	gcctctgagc	tattccagaa	gtagtgagga	ggcttttttg	gaggcctagg	5299											
cttttgcaaa	aagctcccgg	gatggttcga	ccattgaact	gcatcgtcgc	cgtgtcccaa	5359											
aatatgggga	ttggcaagaa	cgggaccta	ccctggccct	cgctcaggaa	cgagttcaag	5419											
tacttccaaa	gaatgaccac	aacctcttca	gtggaaggta	aacagaatct	ggtgattatg	5479											
ggtaggaaaa	cctggttctc	cattcctgag	aagaatcgac	ctttaaaagg	cagaattaat	5539											
atagttctca	cttagaactc	caagaaccaa	ccacgaggag	cctattttct	tgccaaaagt	5599											
ttggatgatg	ccttaagact	tattgaacaa	ccggaattgg	caagtaaagt	agacatggtt	5659											
tggatagtgc	gaggcagttc	tgtttaccag	gaagccatga	atcaaccagg	ccaccttaga</												

WO 02/099092

9/13

PCT/DK02/00388

gagttttcgc cccgaagaac gttttccaat gatgagcact tttaaagtgc tgctatgtgg 6619
cgcggtatta tcccgtattg acgcccggga agagcaactc ggtcgccgca tacactattc 6679
tcagaatgac ttggttgagt actcaccagt cacagaaaag catcttacgg atggcatgac 6739
agtaagagaa ttatgcagtg ctgccataac catgagtgat aacactgcgg ccaacttact 6799
tctgacaacg atcggaggac cgaaggagct aaccgctttt ttgcacaaca tgggggatca 6859
tgtaactcgc ctgtatcgtt gggaaccgga gctgaatgaa gccataccaa acgacgagcg 6919
tgacaccacg atgcctgtag caatggcaac aacgttgccg aaactattaa ctggcgaaact 6979
acttactcta gcttcccggc aacaattaat agactggatg gaggcggata aagttgcagg 7039
accacttctg cgctcggccc ttccggctgg ctgggtttatt gctgataaat ctggagccgg 7099
tgagcgtggg tctcgcggta tcattgcagc actggggcca gatggttaagc cctcccgtat 7159
cgtagttatc tacacgacgg ggagtcaggc aactatggat gaacgaaata gacagatcgc 7219
tgagataggt gcctcactga ttaagcattg gtaactgtca gaccaagttt actcatatat 7279
actttagatt gatttaaaac ttcatTTTTA atttaaaagg atctaggtga agatcctttt 7339
tgataatctc atgaccaaaa tcccttaacg tgagttttcg ttccactgag cgtcagaccc 7399
cgtagaaaaa atcaaaggat cttcttgaga tccttttttt ctgcgcgtaa tctgctgctt 7459
gcaaacaaaa aaaccaccgc taccagcggg gggttggttg ccggatcaag agctaccaac 7519
tctttttccg aaggtaactg gcttcagcag agcgcagata ccaaatactg tccttctagt 7579
gtagccgtag ttaggccacc acttcaagaa ctctgtagca ccgcctacat acctcgctct 7639
gctaatacctg ttaccagtgg ctgctgccag tggcgataag tcgtgtctta ccgggttgga 7699
ctcaagacga tagttaccgg ataaggcgca gcggtcgggc tgaacggggg gttcgtgcac 7759
acagcccagc ttggagcgaa cgacctacac cgaactgaga tacctacagc gtgagctatg 7819
agaaagcgcc acgcttcccc aagggagaaa ggccgacagg tatccggtaa gcggcagggt 7879
cggaacagga gagcgacga gggagcttcc agggggaaac gcctggatc tttatagtcc 7939
tgtcgggttt cgccacctct gacttgagcg tcgatttttg tgatgctcgt caggggggcg 7999
gagcctatgg aaaaacgcca gcaacgcggc ctttttacgg ttccctggcct tttgctggcc 8059
ttttgctcac atggctcgac 8079

<210> 3

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide IC01106

<400> 3

tctgcgctcg aggctgctg gactca

26

<210> 4

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide IC01107

<400> 4

tgctaatacta gagggcccat cccagcagac

30

<210> 5

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide IC0873

<400> 5

tgcagcttat aatggttaca

20

WO 02/099092

10/13

PCT/DK02/00388

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide IC0929

<400> 6
ggcacctatt ggtcttactg 20

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide IC01110

<400> 7
aagtcgtgcg agaccttgtg 20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide IC01111

<400> 8
ccgccaagga atctgtgctg 20

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide IC01112

<400> 9
tacgagcgcc tcagctacaa 20

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide IC01113

<400> 10
gcggaagggtg aattggatgg 20

<210> 11
<211> 20
<212> DNA

WO 02/099092

PCT/DK02/00388

11/13

<213> Artificial Sequence

<220>

<223> oligonucleotide ICO1114

<400> 11

gcctcagggtg cctacatctt

20

<210> 12

<211> 1011

<212> PRT

<213> human

<400> 12

Met	Gly	Ala	Tyr	Ala	Arg	Ala	Ser	Gly	Val	Cys	Ala	Arg	Gly	Cys	Leu
1				5					10					15	
Asp	Ser	Ala	Gly	Pro	Trp	Thr	Met	Ser	Arg	Ala	Leu	Arg	Pro	Pro	Leu
			20					25					30		
Pro	Pro	Leu	Cys	Phe	Phe	Leu	Leu	Leu	Leu	Ala	Ala	Ala	Gly	Ala	Arg
			35				40					45			
Ala	Gly	Gly	Tyr	Glu	Thr	Cys	Pro	Thr	Val	Gln	Pro	Asn	Met	Leu	Asn
	50					55				60					
Val	His	Leu	Leu	Pro	His	Thr	His	Asp	Asp	Val	Gly	Trp	Leu	Lys	Thr
65					70					75				80	
Val	Asp	Gln	Tyr	Phe	Tyr	Gly	Ile	Lys	Asn	Asp	Ile	Gln	His	Ala	Gly
				85					90					95	
Val	Gln	Tyr	Ile	Leu	Asp	Ser	Val	Ile	Ser	Ala	Leu	Leu	Ala	Asp	Pro
			100					105					110		
Thr	Arg	Arg	Phe	Ile	Tyr	Val	Glu	Ile	Ala	Phe	Phe	Ser	Arg	Trp	Trp
			115				120						125		
His	Gln	Gln	Thr	Asn	Ala	Thr	Gln	Glu	Val	Val	Arg	Asp	Leu	Val	Arg
			130			135					140				
Gln	Gly	Arg	Leu	Glu	Phe	Ala	Asn	Gly	Gly	Trp	Val	Met	Asn	Asp	Glu
145					150					155				160	
Ala	Ala	Thr	His	Tyr	Gly	Ala	Ile	Val	Asp	Gln	Met	Thr	Leu	Gly	Leu
				165					170					175	
Arg	Phe	Leu	Glu	Asp	Thr	Phe	Gly	Asn	Asp	Gly	Arg	Pro	Arg	Val	Ala
			180					185					190		
Trp	His	Ile	Asp	Pro	Phe	Gly	His	Ser	Arg	Glu	Gln	Ala	Ser	Leu	Phe
			195				200					205			
Ala	Gln	Met	Gly	Phe	Asp	Gly	Phe	Phe	Phe	Gly	Arg	Leu	Asp	Tyr	Gln
	210					215					220				
Asp	Lys	Trp	Val	Arg	Met	Gln	Lys	Leu	Glu	Met	Glu	Gln	Val	Trp	Arg
					230					235				240	
Ala	Ser	Thr	Ser	Leu	Lys	Pro	Pro	Thr	Ala	Asp	Leu	Phe	Thr	Gly	Val
				245					250					255	
Leu	Pro	Asn	Gly	Tyr	Asn	Pro	Pro	Arg	Asn	Leu	Cys	Trp	Asp	Val	Leu
			260					265					270		
Cys	Val	Asp	Gln	Pro	Leu	Val	Glu	Asp	Pro	Arg	Ser	Pro	Glu	Tyr	Asn
		275					280					285			
Ala	Lys	Glu	Leu	Val	Asp	Tyr	Phe	Leu	Asn	Val	Ala	Thr	Ala	Gln	Gly
	290					295					300				
Arg	Tyr	Tyr	Arg	Thr	Asn	His	Thr	Val	Met	Thr	Met	Gly	Ser	Asp	Phe
	305				310					315				320	
Gln	Tyr	Glu	Asn	Ala	Asn	Met	Trp	Phe	Lys	Asn	Leu	Asp	Lys	Leu	Ile
			325						330					335	
Arg	Leu	Val	Asn	Ala	Gln	Gln	Ala	Lys	Gly	Ser	Ser	Val	His	Val	Leu
			340					345					350		
Tyr	Ser	Thr	Pro	Ala	Cys	Tyr	Leu	Trp	Glu	Leu	Asn	Lys	Ala	Asn	Leu

WO 02/099092

PCT/DK02/00388

12/13

355	360	365
Thr Trp Ser Val Lys His Asp Asp Phe Phe Pro Tyr Ala Asp Gly Pro		
370	375	380
His Gln Phe Trp Thr Gly Tyr Phe Ser Ser Arg Pro Ala Leu Lys Arg		
385	390	395
Tyr Glu Arg Leu Ser Tyr Asn Phe Leu Gln Val Cys Asn Gln Leu Glu		
405	410	415
Ala Leu Val Gly Leu Ala Ala Asn Val Gly Pro Tyr Gly Ser Gly Asp		
420	425	430
Ser Ala Pro Leu Asn Glu Ala Met Ala Val Leu Gln His His Asp Ala		
435	440	445
Val Ser Gly Thr Ser Arg Gln His Val Ala Asn Asp Tyr Ala Arg Gln		
450	455	460
Leu Ala Ala Gly Trp Gly Pro Cys Glu Val Leu Leu Ser Asn Ala Leu		
465	470	475
Ala Arg Leu Arg Gly Phe Lys Asp His Phe Thr Phe Cys Gln Gln Leu		
485	490	495
Asn Ile Ser Ile Cys Pro Leu Ser Gln Thr Ala Ala Arg Phe Gln Val		
500	505	510
Ile Val Tyr Asn Pro Leu Gly Arg Lys Val Asn Trp Met Val Arg Leu		
515	520	525
Pro Val Ser Glu Gly Val Phe Val Val Lys Asp Pro Asn Gly Arg Thr		
530	535	540
Val Pro Ser Asp Val Val Ile Phe Pro Ser Ser Asp Ser Gln Ala His		
545	550	555
Pro Pro Glu Leu Leu Phe Ser Ala Ser Leu Pro Ala Leu Gly Phe Ser		
565	570	575
Thr Tyr Ser Val Ala Gln Val Pro Arg Trp Lys Pro Gln Ala Arg Ala		
580	585	590
Pro Gln Pro Ile Pro Arg Arg Ser Trp Ser Pro Ala Leu Thr Ile Glu		
595	600	605
Asn Glu His Ile Arg Ala Thr Phe Asp Pro Asp Thr Gly Leu Leu Met		
610	615	620
Glu Ile Met Asn Met Asn Gln Gln Leu Leu Leu Pro Val Arg Gln Thr		
625	630	635
Phe Phe Trp Tyr Asn Ala Ser Ile Gly Asp Asn Glu Ser Asp Gln Ala		
645	650	655
Ser Gly Ala Tyr Ile Phe Arg Pro Asn Gln Gln Lys Pro Leu Pro Val		
660	665	670
Ser Arg Trp Ala Gln Ile His Leu Val Lys Thr Pro Leu Val Gln Glu		
675	680	685
Val His Gln Asn Phe Ser Ala Trp Cys Ser Gln Val Val Arg Leu Tyr		
690	695	700
Pro Gly Gln Arg His Leu Glu Leu Glu Trp Ser Val Gly Pro Ile Pro		
705	710	715
Val Gly Asp Thr Trp Gly Lys Glu Val Ile Ser Arg Phe Asp Thr Pro		
725	730	735
Leu Glu Thr Lys Gly Arg Phe Tyr Thr Asp Ser Asn Gly Arg Glu Ile		
740	745	750
Leu Glu Arg Arg Arg Asp Tyr Arg Pro Thr Trp Lys Leu Asn Gln Thr		
755	760	765
Glu Pro Val Ala Gly Asn Tyr Tyr Pro Val Asn Thr Arg Ile Tyr Ile		
770	775	780
Thr Asp Gly Asn Met Gln Leu Thr Val Leu Thr Asp Arg Ser Gln Gly		
785	790	795
Gly Ser Ser Leu Arg Asp Gly Ser Leu Glu Leu Met Val His Arg Arg		
805	810	815
Leu Leu Lys Asp Asp Gly Arg Gly Val Ser Glu Pro Leu Met Glu Asn		
820	825	830

WO 02/099092

PCT/DK02/00388

13/13

Gly Ser Gly Ala Trp Val Arg Gly Arg His Leu Val Leu Leu Asp Thr
835 840 845
Ala Gln Ala Ala Ala Ala Gly His Arg Leu Leu Ala Glu Gln Glu Val
850 855 860
Leu Ala Pro Gln Val Val Leu Ala Pro Gly Gly Gly Ala Ala Tyr Asn
865 870 875 880
Leu Gly Ala Pro Pro Arg Thr Gln Phe Ser Gly Leu Arg Arg Asp Leu
885 890 895
Pro Pro Ser Val His Leu Leu Thr Leu Ala Ser Trp Gly Pro Glu Met
900 905 910
Val Leu Leu Arg Leu Glu His Gln Phe Ala Val Gly Glu Asp Ser Gly
915 920 925
Arg Asn Leu Ser Ala Pro Val Thr Leu Asn Leu Arg Asp Leu Phe Ser
930 935 940
Thr Phe Thr Ile Thr Arg Leu Gln Glu Thr Thr Leu Val Ala Asn Gln
945 950 955 960
Leu Arg Glu Ala Ala Ser Arg Leu Lys Trp Thr Thr Asn Thr Gly Pro
965 970 975
Thr Pro His Gln Thr Pro Tyr Gln Leu Asp Pro Ala Asn Ile Thr Leu
980 985 990
Glu Pro Met Glu Ile Arg Thr Phe Leu Ala Ser Val Gln Trp Lys Glu
995 1000 1005
Val Asp Gly
1010

